

United States
Department of
Agriculture

Marketing and
Regulatory
Programs

Animal and
Plant Health
Inspection Service

Cooperating State
Departments of
Agriculture

Emerald Ash Borer Program Manual

Agrilus planipennis (Fairmaire)



Emerald Ash Borer Program Manual, *Agrilus planipennis* (Fairmaire), ver. 1.1

Prepared by:

James H. Buck, Ph.D.
USDA APHIS PPQ EAB National Program Analyst
Emerald Ash Borer Program

Susan Frappier
USDA APHIS PPQ Supervisory Officer
Emerald Ash Borer Program

ACKNOWLEDGEMENTS

Considerable assistance, guidance, and review of sections in this Program Manual was contributed by: Philip Bell, Paul Chaloux, Kevin Dodds, Jody Feliciano, Joel Floyd, Paula Henstridge, Craig Kellogg, Sharon Lucik, Deborah McPartlan, Michelle Mikula, Elizabeth Pentico, Michael Stefan, Ken Witt, and Dr. James Zablotny. Camille Chapman also assisted with formatting.

Cite this report as follows: USDA–APHIS. 2009. *Emerald Ash Borer Program Manual, Agrilus planipennis* (Fairmaire) USDA–APHIS–PPQ–Emergency and Domestic Programs–Emergency Planning, Riverdale, Maryland.

Cover Illustration: Joel Floyd

CONTENTS

SECTION	PAGE
ACKNOWLEDGEMENTS	ii
CONTENTS	iii
LIST OF TABLES AND FIGURES	vi
1) INTRODUCTION	
a) Purpose	1.1
b) Disclaimers	1.1
c) Program Safety	1.1
d) Support for Program Decision-Making	1.2
2) PEST INFORMATION	
a) Systematic Placement	2.1
b) Background Information	2.1
c) Historical Information	2.1
d) Economic Impact	2.3
e) Host Range	2.4
f) Geographic Distribution	2.4
g) Biology	2.5
h) Life Cycle	2.5
i) Development	2.7
3) IDENTIFICATION	
a) Importance	3.1
b) Authorities	3.1
c) Identification	3.1
d) Similar Species	3.4
e) Collection and Preparation of Specimens	3.8
f) Screening for Suspect Buprestidae and Specimen Submission	3.8
4) RESPONSE PROCEDURES	
a) Introduction	4.1
b) Consultation	4.1
c) New State Detection	4.1
d) Identification by APHIS Identifier	4.1
e) Confirmation by SEL	4.1
f) Public Declaration and SPRO Memo	4.1
g) Radial Delimiting Survey	4.2
h) Analysis of Survey Data	4.2
i) Community Management Plan	4.2
j) Biological Control	4.2

k) Eradication	4.2
l) Currently	4.3
m) Long Range Management Options	4.3
5) SURVEY PROCEDURES	
a) Introduction	5.1
b) Trace Back and Trace Forward Investigations	5.1
c) General Detection Surveys	5.1
d) Visual Survey	5.2
e) Symptoms of EAB Infestations	5.3
f) Delimiting Sampling	5.5
g) Destructive Survey	5.5
h) Monitoring Survey	5.6
i) Survey Sample Collection	5.6
j) Quality Control for Survey Activities	5.6
6) REGULATORY PROCEDURES	
a) Instructions to Officers	6.1
b) Authorities	6.1
c) Regulating the Spread of EAB	6.1
d) Regulated Articles	6.2
e) Approved Regulatory Treatments	6.2
f) Quarantine Boundaries	6.3
g) Regulatory Management of Outlying Infestations	6.3
h) Regulated Establishments	6.4
i) Principal Regulatory Activities	6.4
j) Investigations and Violations	6.5
k) Regulatory Records	6.6
l) Quality Control	6.6
m) Outreach	6.6
n) Special Regulatory Operations	6.6
o) Issuing an Emergency Action Notification	6.7
p) Removing Areas from Quarantine	6.7
q) Regulatory Forms	6.7
7) MANAGEMENT PROCEDURES	
a) Overview	7.1
b) Community Preparedness Planning	7.1
c) Biological Control	7.2
d) Chemical Control	7.3
e) Eradication Strategy	7.4
f) Management Records	7.4
g) Monitoring	7.4
8) ENVIRONMENTAL COMPLIANCE	
a) Overview	8.1
b) Disclaimer	8.1

c) Protected Species	8.1
d) Environmental Assessment	8.1
9) PUBLIC OUTREACH	
a) Introduction	9.1
b) General Outreach Objectives	9.1
c) General Outreach Activities and Initiatives	9.2
d) Outreach Material	9.3
e) Telephone Hotline	9.3
f) Web Site	9.3
g) Public Meetings or Informational Open Houses	9.3
h) Media Relations	9.4
10) REFERENCES	10.1
11) DEFINITIONS	11.1
12) APPENDICES	
Appendix A - <i>Agrilus planipennis</i> (Fairmaire) Screening Aid	A-1
Appendix B - Emerald Ash Borer Larval Screening Guide	B-1
Appendix C – Emerald Ash Borer Program Contacts	C-1
Appendix D – Information Packet	D-1
Appendix E – Regulatory Operations Forms	E-1
Appendix F – Sources to Find Regulatory Concerns	F-1
Appendix G – PPQ Form 518 (Report of Violation)	G-1
Appendix H – PPQ Form 523 (Emergency Action Notification)	H-1
Appendix I – PPQ Form 530 (Limited Permit)	I-1
Appendix J – PPQ Form 540 (Certificate)	J-1
Appendix K - Passive dispersal pathways of emerald ash borer, <i>Agrilus planipennis</i> (Fairmaire), (Coleoptera: Buprestidae) identified by USDA APHIS PPQ EAB Program	K-1

TABLES AND FIGURES		PAGE
Figure 2.1	Emerald ash borer life cycle	2.5
Figure 3.1	<i>Agrilus planipennis</i> egg on bark	3.1
Figure 3.2	<i>A. planipennis</i> egg cluster	3.1
Figure 3.3	Unhatched <i>Agrilus planipennis</i> neonate on bark	3.2
Figure 3.4	Three instars of <i>A. planipennis</i> larvae	3.2
Figure 3.5	Anterior abdominal segments	3.2
Figure 3.6	Posterior abdominal segments	3.3
Figure 3.7	<i>Agrilus planipennis</i> prepupal stage	3.3
Figure 3.8	Pupal ventral and dorsal views	3.3
Figure 3.9	<i>Agrilus planipennis</i> adult	3.4
Figure 3.10	<i>Agrilus anxius</i> and <i>A. bilineatus</i> adults	3.5
Figure 3.11	<i>Agrilus bilineatus</i> and <i>A. cyanescens</i> adults	3.6
Figure 3.12	<i>Agrilus masculinus</i> and <i>A. vittaticollis</i> adults	3.7
Figure 3.13	Emerald Ash Borer: Specimen Processing and Communication Protocol	3.10
Figure 4.1	New Detection Decision Framework	4.3
Figure 5.1	Symptoms of EAB on <i>Fraxinus spp.</i>	5.3
Figure 5.2	Symptoms of EAB on <i>Fraxinus spp.</i>	5.4
Table 6.1	Interstate movement of regulated articles from quarantined areas	6.2

Purpose

These New Pest Response Guidelines contain information to guide a management program for the emerald ash borer (EAB) beetle, *Agrilus planipennis* (Fairmaire), (Coleoptera: Buprestidae).

The guidelines are intended to assist Plant Protection and Quarantine field operations and states in implementing specific action plans to manage infestations and prevent the spread of EAB to other locations. This information provides strategies for detection and response to an infestation of EAB by presenting available information for implementing general and delimiting surveys, identification, regulatory, management, or containment procedures. Specific program activity should be based on information available at that time.

Plant Protection and Quarantine (PPQ) develops guidelines through discussion, consultation, or agreement with other Animal and Plant Health Inspection Service (APHIS) staff, State Plant Regulatory Officials (SPRO), U.S. Forest Service, tribal governments, and other state and federal agencies and cooperators directly involved in EAB management. The APHIS Emergency Response Programs Manual and other pest national response guidelines may be found at:

http://www.aphis.usda.gov/import_export/plants/manuals/emergency/index.shtml

Disclaimers

Document comprehensiveness: This document is not intended to be complete and exhaustive, but a foundation based on literature available to assist future EAB control efforts. Few publications were available at the time of this writing and not all specialists and members of the research community were consulted for their advice. As ongoing research and program efforts provide new information, the direction and strategies of the guidelines may change to reflect the best management practices for the control of EAB. For the latest updates on this pest, it is recommended to conduct periodic literature searches on the web and in other archives.

Commercial Suppliers or Products: References to commercial suppliers or products should not be construed as an endorsement of the company or product by the U.S. Department of Agriculture.

Program Safety

Safety of the public and program personnel has priority in preprogram planning and training, and throughout operations. Safety officers and supervisors must enforce on-the-job safety procedures.

**Support for
Program Decision-
Making**

The USDA APHIS PPQ Center for Plant Health, Science and Technology (CPHST) provides technical support, in consultation with other scientists, to emergency pest response program directors concerning risk assessments, survey methods, management strategies, regulatory treatments, and other aspects of the pest response program.

Systematic Placement

Phylum: Arthropoda
 Class: Insecta
 Order: Coleoptera
 Family: Buprestidae
 Genus: *Agrilus*
 Species: *planipennis*
 Entomological Author: Fairmaire, Léon Marc Herminie

Approved Name: *Agrilus planipennis* (Fairmaire)

Synonyms: *Agrilus marcopoli* (Obenberger 1930)
Agrilus marcopoli ulmi (Kurosawa 1956)
Agrilus feretrius (Obenberger 1936)

Common Names: emerald ash borer (English)
 agrile du frêne (French)
 изумрудная ясеневая златка (Russian)

Background Information

Emerald ash borer (EAB) is a non-native wood-boring pest of North American ash trees. This devastating pest was first found in 2002 in North America where it was discovered in southeastern Michigan and adjacent areas in Windsor, Ontario, Canada. It is thought to have been introduced in the 1990's on solid wood packing material originating from Asia.

This extremely destructive beetle poses an enormous threat to all of North America's ash resources. Unlike many other wood boring beetles, EAB aggressively kills healthy and stressed trees; many dying within two to three years after becoming infested. Currently, EAB has no known natural enemies in North America and no effective control options. If it is not contained or its effects mitigated, this pest will continue to infest and kill all species of trees in the genus *Fraxinus*. The impact on ash in North America has been compared to the effects of chestnut blight and Dutch elm disease, which devastated rural and urban forests in the 20th century.

Historical Information

Biological information on EAB is scarce. Prior to 2002, only two short papers occurred in the literature: Chinese Academy of Science (1986) and Yu (1992). These papers include brief morphology, biology, host range, and symptoms of infestation.

In 2002, shortly after EAB was confirmed as the cause of significant ash tree mortality observed in Detroit, Michigan, five counties were found to be infested and were placed under quarantine. Since that time, APHIS, U.S. Forest Service, state, and local cooperators have conducted

survey, control, and eradication activities. Efforts have included imposing quarantines, conducting surveys, delimiting areas around confirmed infested sites, removing ash trees, and developing information which will support management efforts. Lack of effective survey and control technology has made containment efforts challenging.

Intensive visual survey efforts in 2003 expanded this area by 12 additional counties in Michigan. After intensive survey efforts in northern Ohio, EAB was discovered there in three counties. Additionally, due to a 2002 Michigan quarantine violation involving nursery stock, control actions were implemented in Maryland and Virginia in 2003.

Survey methods were improved in 2004 through the implementation of trap tree survey, a survey method more focused than visual survey. This survey method, in combination with the visual survey methods, resulted in the discovery of 27 additional counties in 2004: 20 in Michigan, five in Ohio, and two in Indiana discovered in April and May.

In 2005, one additional new detection was made in Indiana, five in Ohio, and 16 in Michigan which accounted for the 22 new county finds that year.

Illinois was found to be infested in 2006; the first infested county was found in June, the second in July. This was in addition to 12 additional new county finds in Michigan, 11 in Ohio, and seven in Indiana, for a total of 32 new county finds for 2006. In addition, Maryland reported a re-infestation of a site that underwent eradication actions in 2004.

Michigan found EAB in seven additional counties in 2007. Indiana discovered five more infested counties and Ohio saw an increase of 11 more counties this same year. Two more counties were found to have EAB in Illinois and the state of Pennsylvania reported their initial infestation of two counties.

To date, 2008 currently has 24 additional new reports in Ohio, Indiana, Michigan, Illinois, Pennsylvania, Maryland, and new infestations in singular counties of Missouri and Virginia.

The first state and federal quarantines began in 2002 with the original six infested Michigan counties. This area was expanded in 2003 to a total of 13 counties and two portions of counties in Michigan, four portions of counties in Ohio, and one portion of a county in Maryland. At the end of 2004, 20 entire counties in Michigan were considered generally infested and quarantined; portions of 16 additional counties in

Michigan, four counties in Ohio, and two counties in Indiana were quarantined due to small spot infestations caused by the movement of firewood, nursery stock, or timber. In 2005, the quarantine expanded to 21 entire counties and 25 partial counties in Michigan, four entire counties in Indiana, and one county and ten partial counties in Ohio.

By December, 2006, the entire Lower Peninsula of Michigan (68 counties) as well as one partial county in the Upper Peninsula, the entire states of Ohio, Indiana, and Illinois and a county in Maryland were federally quarantined to prevent the spread of this destructive pest.

On August 8, 2007 the federal quarantine was expanded to include four counties in Pennsylvania after discovery of an initial infestation in that state.

Wayne County incurred the initial infestation in Missouri and was added to the federal quarantine on August 8, 2008. Four counties (Ozaukee, Washington, Fond du Lac and Sheboygan) in Wisconsin were federally quarantined on August 12, 2008. On December 4, 2008 the counties of Delta, Mackinac, and Schoolcraft in the Upper Peninsula of Michigan were added to the Federal Quarantine.

Economic Impact

The eastern United States produces nearly 114 million board feet of ash saw timber with a value of \$25.1 billion (McPartlan et al. 2006). White, black, and green ash make up over 7 percent of all hardwood species and 5.5 percent of all tree species in the northeastern United States and eastern Canada. The wood is used for a variety of applications including tool handles, baseball bats, furniture, cabinetry, solid wood packing materials, pulp, and paper. The continued spread of this pest threatens these resources and may permanently alter landscape ecosystems of the Midwest, which consists of up to 20 to 40 percent ash in some areas.

There is potential for extensive negative economic effects if this wood-borer were to become widespread in the United States. If left unchecked, EAB will continue to infest and destroy ash trees, resulting in the losses of millions of dollars to the lumber and nursery industries as well as urban communities. Preliminary findings by U.S. Forest Service estimate that EAB's potential impact to the national urban landscape is a potential loss of between 0.5 to 2 percent of the total leaf area (30-90 million trees) and a value loss of between \$20-60 billion (McPartlan et al. 2006).

Infested states may experience significant economic losses in forest products if EAB spreads from the currently quarantined area into the forests of the eastern United States. Quarantines imposed by state and

federal agencies have negative impacts on the nursery, landscaping, timber, recreation, and tourism industries which are economically important to the region.

In addition to its value to the timber industry and the forest ecosystem, ash is one of the most popular landscape trees because of its wide ecological amplitude and resistance to other pests. It is the most commonly planted tree in new residential and commercial developments.

In an initial economic analysis of EAB, the U.S. Forest Service estimated that EAB, if not contained and eradicated, could cause approximately \$7 billion in additional costs to state and local governments and landowners to remove and replace dead and dying ash trees in urban and suburban areas over the next 25 years. Nationwide, the nursery industry produces an estimated 2 million ash trees each year. With median approximate values ranging from \$50 to \$70 per tree, the ash nursery stock crop is worth between \$100 and \$140 million annually (McPartlan et al. 2006).

Host Range

In North America, EAB is capable of infesting all ash trees in the genus *Fraxinus*, including green ash (*F. pennsylvanica*), white ash (*F. americana*), black ash (*F. nigra*), pumpkin ash, (*F. profunda*), blue ash (*F. quadrangulata*), and other native species in this same genus (MacFarlane and Meyer, 2005). An approximate natural range map of all ash species in North America can be found in Appendix D.

Chinese reports indicate that the species *F. chinensis* var. *chinensis*, *F. chinensis* var. *rhynchophylla*, and *F. mandshurica*. (Chinese Academy of Science 1986, Yu 1992) are native hosts in Asia.

In Japan, the host range includes Manchurian ash, (*Fraxinus mandshurica* var. *japonica*), Manchurian walnut (*Juglans mandshurica* var. *sieboldiana*, and var. *sachalinensis*), Japanese wingnut (*Pterocarya rhoifolia*) and Japanese elm (*Ulmus davidiana* var. *japonica*), (Akiyama and Ohmomo 1997, Sugiura 1999).

There is an isolated reference to privet (*Ligustrum* spp.) being suitable for 1st stage larval development in a laboratory setting (Cappaert et al. 2005).

Geographic Distribution

Native distribution of EAB in Asia includes several provinces of China (Liaoning, Jilin, Heilongjiang, Inner Mongolia, Hebei, and Shandong), Korea, Japan, Taiwan, and a small area in adjacent Russia and Mongolia

(US Forest Service 2008).

EAB is now considered established in portions of the United States and Canada. The most current map depicting the emerald ash borer infestation can be found at:

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/index.shtml

Biology

Current research suggests that EAB can complete either a one or two year life cycle. It has been observed that low density populations on vigorous ash trees tend to support a two year life cycle while stressed trees with higher beetle population densities tend to support a one year life cycle (Cappaert et al. 2005). Effects on life cycle due to latitudinal and altitudinal variation are unknown at this time.

Life Cycle




Figure 2.1 Emerald ash borer life cycle.

Eggs: This life stage is extremely difficult to visually detect during survey. Oviposition sites are likely to be correlated with the direction of sunlight, with the most eggs being laid on the southwestern side of the tree (Timms et al. 2006). Recent research indicates that females may prefer to lay eggs in areas where the bark is rough, cracked, or rippled, as may be found near branch crotches. The female may lay 1-23 eggs at

a time, with one being the norm. Each female can lay 60-90 eggs in their lifetime, depositing them individually on the bark along the trunk and portions of the major branches. The eggs typically hatch in 7-10 days (Yu 1992).

Larva: Minute larvae bore through the bark and into the cambium where they feed on the phloem from mid-June to mid-October. There are four stages to larval development (Cappaert et al. 2005). As they feed, the larvae create long serpentine galleries (20-30 cm) filled with frass, which enlarge in width as they grow. Larvae continue development into fall. Facultative diapause occurs after fourth instar larvae enter the sapwood or outer bark and excavate a pupal chamber where they overwinter as prepupae. Larvae too immature to prepupate spend the winter in the larval gallery and complete development the following spring. Larval galleries are typically 20-30 cm long.

Pupa: Pupal development is variable according to humidity and temperature. Pupation may begin upon the accumulation of 100-150 growing degree days and, in ideal laboratory conditions, pupal development typically takes approximately 4 weeks. After pupae transform into adults, the beetle takes 1 to 2 weeks before it emerges through D-shaped exit holes 3-4 mm wide. More research is necessary to fully understand pupal development.

Adult: Newly formed adults typically remain in the pupal chamber for 1-2 weeks after pupation is complete. Initial adult emergence predictively occurs when an accumulation of 400-500 growing degree days is achieved. Evidence of an adult emergence appears in the form of a D-shaped exit hole 3-4 mm in diameter. Peak activity for adults is predicted to occur at approximately 1,000 growing degree days (McCullough and Siegert 2006). Adults are capable of immediate flight and, in laboratory conditions, adults mate shortly after emergence. An approximate 3 week period of maturation feeding occurs before oviposition. The adults feed on ash foliage, causing minimal damage. Adults may feign death when frightened or disturbed. They are phototactic and thermotactic and most active on warm, cloudless, windless days. Oviposition begins 7-9 days after the initial mating (Yu 1992). Average longevity for adult males is 13 days; average longevity for females is 22 days (Bauer et al. 2004, Lyons et al. 2004, Poland and McCullough 2006). *Agrilus planipennis* beetles are strong fliers, with females flying twice as far as males and mated females flying twice as far as unmated females. Tethered flight in laboratory conditions suggests that a mated female may fly more than 20 km (Taylor et al. 2006). Females may mate as many as 3 times with mating lasting 20-90 minutes (average of 60 min.).

Development

Many environmental factors can influence the development of insects, the timing of their biological events, and the dynamics of their populations. Among these factors are host availability, population densities, photoperiod, and weather. Temperature and moisture, because they are so critical to biochemical reactions, are universal influences on egg, larval, pupal and adult development in insects. Current research suggests initial adult emergence occurs when cumulative growing degree days reach 400-500 degree days with peak activity occurring at approximately 1000 growing degree days. Temporal variation for these occurrences may exceed one month when considering latitudinal differences in the current quarantine area.

More research is necessary to further understand EAB biology, including developmental thresholds on its life cycle.

Importance Accurate identification of the pest is pivotal to assessing its potential risk, developing a survey strategy, and deciding the level and manner of control and/or management.

Authorities For new detections in a state or county, the USDA APHIS PPQ National Identification Service must positively identify the suspect pest as *Agrilus planipennis* before consideration of any containment, control, regulatory, or eradication activities.

Identification Some pre-identification and screening can be performed by field personnel assigned to the program if training is provided. A description of EAB, *Agrilus planipennis*, including distinctive features that separate it from native species that resemble it, with pictures, occurs below.

Description of the species, *Agrilus planipennis*

Eggs

Eggs are 1.0 mm long x 0.6 mm wide and oblate. They are creamy white turning to amber before hatching with a reductus extending radially toward the edges. Extremely difficult to observe with the naked eye.



Figure 3.1 *Agrilus planipennis* egg on bark. Photo: Houping Liu, www.forestryimages.org



Figure 3.2 *A. planipennis* egg cluster. Photo: David Cappaert, Michigan State University



Figure 3.3 Unhatched *Agrilus planipennis* neonate on bark. Photo: Houping Liu, www.forestryimages.org

Larvae

Larvae are creamy white, and dorso-ventrally flattened. They are 26-32 mm long when fully mature in the fourth instar. The small, brown head is mostly retracted into the prothorax with only the mouthparts remaining visibly externally. The prothorax is enlarged with the mesothorax and metathorax more narrow. The mesothorax and each of the first eight abdominal segments have a pair of spiracles.

There are ten abdominal segments. The first abdominal segment shape varies considerably but can be distinctly rectangular to trapezoidal shaped. Segments A2 through A6 are somewhat trapezoidal with protruding flattened



Figure 3.4 Three instars of *A. planipennis* larvae. Photo: David Cappaert, Michigan State University



Figure 3.5 Anterior abdominal segments. Photo: James Zabltny, USDA APHIS PPQ

3. Identification

Emerald Ash Borer

lobes. A7 is strongly bell shaped and A8 is somewhat trapezoidal in shape. The last abdominal segment (A10) contains a pair of urogomphi. Emerald ash borer prepupae are more difficult to identify, being subtly different than less mature larvae, but A8 is still bell shaped.




Figure 3.6 Posterior abdominal segments. Photo: James Zablony, USDA APHIS PPQ

A larval screening guide comparing *Agrilus planipennis* to *A. anxius* may be found in Appendix L.

Pupae

Pupae are creamy white and 10-14 mm long. The antennae extend dorsally to the base of the wing buds. The posterior abdominal segments are slightly curved ventrally.



Fig 3.7 *Agrilus planipennis* prepupal stage



Figure 3.8 Pupal ventral and dorsal views Photo: Deborah Miller, USFS (NC)

Adults

Adults have metallic (brassy or golden) green pronotum, with darker, metallic emerald green elytra and abdominal sternites. They may reach a



Figure 3.9 *Agrilus planipennis* adult. Photo: David Cappaert, Michigan State University

length of 8.5-13.5 mm long and 3.1-3.4 mm wide. The cuneiform body is narrow and elongate. The abdominal tergites are metallic coppery red. The head is flat with the vertex shield shaped. The compound eyes are obscure-aeueous and kidney shaped. The rectangular prothorax is slightly wider than the head, but the same width as the elytra. The anterior margin of the elytra is raised, forming a transverse ridge; the surface is covered with punctures. The elytra's posterior margins are round and obtuse with small denticles on the edge. There is an emarginate pygidial spine.

Similar Species

Agrilus is one of the largest genera in the world with almost 3,000 described species. Species in this genus are difficult to identify because of structural coloration (*i.e.*, *Agrilus bilineatus* in Fig. 3.10 and Fig. 3.11) and subtle morphological differences.

Agrilus also is a very diverse genus with 171 known species in North America. North American species most similar to *Agrilus planipennis* are *A. anxius* (Weber) (Fig. 3.10), *A. bilineatus* (Weber) (Fig. 3.10 and Fig. 3.11), *Agrilus cyanescens* Ratzeburg (Fig. 3.11), *A. masculinus* Horn (3.12), *A. obsoletoguttatus* Gory, *A. subcinctus* Gory, and *A. vittaticollis* (Randall) (Fig. 3.12).



Agrilus anxius (Weber)



Agrilus bilineatus (Weber)

Figure 3.10 *Agrilus anxius* and *A. bilineatus* adults



Agrilus bilineatus (Weber)



Agrilus cyanescens Ratzeburg

Figure 3.11 *Agrilus bilineatus* and *A. cyanescens* adults



Agrilus masculinus Horn



Agrilus vittaticollis (Randall)

Figure 3.12 *Agrilus masculinus* and *A. vittaticollis* adults

The only other *Agrilus* species in the Midwest besides *Agrilus planipennis* known to occur on *Fraxinus* spp. is *Agrilus subcinctus*. A major behavioral difference between *A. subcinctus* and *A. planipennis* is that *A. subcinctus* is a twig borer and prefers smaller branches for oviposition.

Collection and Preparation of Specimens

Collect as many specimens as possible of the pest for identification. Do not mix samples. Be sure to separate larvae into vials by tree or location. Use of entomological forceps is recommended in order to minimize damage to the specimen. Collect additional information as noted:

- Fill out PPQ 391.
- Gather the following information about the tree:
 - State
 - County
 - Date
 - GPS Coordinates
 - Tree Diameter at Breast Height (DBH)

Prepare specimens according to the following protocols.

- ▣ Gather larvae/pupae from the same tree into the same vial. Do not insert larvae into alcohol at this time.
- ▣ Label the vial with the naming protocol: Year-collector initials-Month-Date (for example: 06-PDB-10-29)
- ▣ Larvae should be killed in boiling water, allowed to cool, and placed in 70% ethyl alcohol or 70% isopropyl alcohol prior to shipping.
- ▣ If there are too many larvae/pupae, extras may be placed in alcohol in larger container (labeled completely)
- ▣ Ship vials in a well-padded box (be aware of possible restrictions on shipping alcohol by air carriers).

Large specimens or small specimens that have been crowded into one vial should be transferred to fresh alcohol within a day or two to reduce the danger of diluting the alcohol with insect body fluids. If the alcohol becomes too diluted, the specimens may begin to decompose. It is recommended to place vials in a vapor lock bag to prevent leakage.

Screening for Suspect Buprestidae and Specimen Submission

Any suspect Buprestidae adult or suspect EAB specimen collected from a trap in a **non-quarantined** state should be placed in a vial with 70% ethanol and delivered to the State Plant Health Director or APHIS representative to be packaged and shipped to Dr. James Zablotny along with a completed "Specimens for Determination" PPQ form 391. Be

3. Identification

Emerald Ash Borer

sure to include any survey record number and/or GPS coordinates on the PPQ form 391 so identified specimens can be linked to survey records. Additional data should also include:

1. EAB detected by: Program related personnel or non-Program related personnel
2. EAB detected in: Purple trap - Detection tree - Destructive Sampling- Non-targeted ash tree
3. Is this a single occurrence or are other symptoms and life stages present? Please specify _____.

Dr. James Zablotny
USDA, APHIS, PPQ
11200 Metro Airport Center Drive, Suite 140
Romulus, MI 48174

Phone: 734-942-9005
e-mail: james.e.zablotny@aphis.usda.gov

Dr. Zablotny will make a determination and send specimens to the Systematic Entomology Laboratory (SEL) if necessary for initial state detection confirmation.

Any suspect Buprestidae adult or suspect EAB specimen collected from a trap in a **quarantined** state should be placed in a vial with 70% ethanol and delivered to the State Plant Health Director or APHIS representative to be packaged and shipped to Dr. Bobby Brown along with a completed "Specimens for Determination" PPQ form 391. Be sure to include any survey record number and/or GPS coordinates on the PPQ form 391 so identified specimens can be linked to survey records. Additional data should also include:

1. EAB detected by: Program related personnel or non-Program related personnel
2. EAB detected in: Purple trap - Detection tree - Destructive Sampling- Non-targeted ash tree
3. Is this a single occurrence or are other symptoms and life stages present? Please specify _____.

Dr. Bobby Brown
USDA, APHIS, PPQ
901 W. State Street
Smith Hall, Purdue University

3. Identification

Emerald Ash Borer

West Lafayette, IN 47907-2089

Phone: 765-496-9673

e-mail: robert.c.brown@aphis.usda.gov




Figure 3.13 Emerald Ash Borer: Specimen Processing and Communication Protocol.

Introduction	Emerald Ash Borer (EAB) is a difficult pest to detect during early stages of infestation. Following introduction, visual signs or symptoms of infestation are often not observed for several years. When obvious visual signs of infestation (e.g., D-shaped holes, bark splits, crown dieback) are observed the possibility of eradication is remote. The EAB Decision Framework is an overview of what to expect and procedures to generally follow when a state's initial infestation is detected.
Consultation	It is imperative for the State Plant Regulatory Official (SPRO), State Plant Health Director (SPHD), and local officials to understand the initial steps to be taken when a new state detection is confirmed. Early in the identification and confirmation cycle discussions regarding possible regulatory actions and operational activities should be held with the National Program Coordinator and the appropriate Regional Program Manager in order to coordinate all involved parties on these issues.
New State Detection	Development of a lure-baited sticky trap has provided a new method of detecting the presence of EAB. In addition, outreach efforts have educated the general public on the identification of signs and symptoms of EAB infestation. New state finds may be identified in a variety of ways: <ol style="list-style-type: none">Caught in a baited prism trapReported by local government official, arborists, industry, general public, etc.
Identification by APHIS Identifier	Collected suspect is submitted to the APHIS Identifier along with PPQ Form 391 (See Fig. 3.13). A suspect identified as EAB is then forwarded to the USDA Systematic Entomology Laboratory (SEL) for confirmation. Negative findings will be communicated to the submitter.
Confirmation by SEL	Notification of SEL confirmation of EAB is forwarded to Domestic Diagnostics Coordinator with USDA APHIS PPQ National Identification Service (NIS). NIS then notifies Emergency and Domestic Programs Staff who in turn notify the respective SPRO and SPHD and other involved parties.
Public Declaration and SPRO Memo	Making the first public announcement of the new find is an option available to the affected state. A discussion is held with the National EAB Program Coordinator, Regional Program Manager and the SPHD and SPRO in the affected state to determine if the state would like to provide first notice and to discuss the timing of state and federal notices of the new find. PPQ's Deputy Administrator confers with the senior state agriculture official prior to release of the SPRO memo.

- Radial Delimiting Survey** An initial visual delimiting survey of infestations accompanied by trees exhibiting obvious signs (*e.g.*, D-shaped exit holes, epicormic sprouting, crown dieback, serpentine galleries, bark splits, woodpecker damage) should be conducted immediately. This initial survey is intended to provide a preliminary sense of the size and scope of the infestation. A more intensive follow-up survey strategy will be developed in consultation with the EAB Program Management.
- Survey Data** Compilation of the number of infested trees, larvae, prepupae, and adults and plotting the location where each was found on a map will aid in indicating epicenter and extent of the new infestation. This data should be input to the [Survey Data Sheet](#) for ISIS. Inspecting in and near high risk facilities such as campgrounds, sawmills, new construction (nursery stock) may aid in determining the origin of the infestation.
- Community Management Plan** Since cost effective control options to eradicate or prevent dispersal are not available at this time, it is recommended that authorities plan and prepare for an infestation of emerald ash borer.
- Examples of preparedness or response plans for states may be found at:
- <http://www.emeraldashborer.info/communityplan.cfm>
- Biological Control** An EAB rearing facility for biocontrol agents has been established at the APHIS facility in Brighton, Michigan. Tools and techniques for rearing and releasing natural enemies of EAB will be refined at this location with the goal of future distribution of parasitoids to other states.
- Three parasitoid stingless wasps are currently under evaluation by APHIS. These parasitoids are known to attack EAB in its native range in China.
- Small-scale trial releases were performed in 2007, 2008 and, 2009. Release sites will be monitored for establishment of the natural enemies and evaluated for their potential to control and slow the dispersal of EAB in other states.
- Eradication** The Emerald Ash Borer Program has transitioned from an eradication program to an integrated pest management (IPM) program. Effective and cost efficient control and eradication technologies are not currently available. In the future additional tools and strategies may become available to suppress the dispersal of this pest. Program partners are conducting extensive research to provide additional IPM tools and methodologies.

Currently The EAB National Program established a pest detection survey which utilizes baited panel traps in 48 states. Purple panel traps are placed in high-risk sites and in a band around the perimeter of the generally infested area. Work is currently underway to assess data from past surveys in order to improve and enhance detection survey work in upcoming years.

Long Range Management Options Continued regulation and outreach are currently the most effective tools to prevent the dispersal of EAB. In the near future, biocontrol agents may be available to reduce and manage EAB populations. Ongoing research into promising new tools will help to prevent dispersal through development of more effective traps, more attractive lure, and one or more effective pesticides.




Figure 4.1 New Detection Decision Framework (adapted from Biosecurity New Zealand, Ministry of Agriculture and Forestry, Wellington, New Zealand).

Introduction Surveys support the program by providing information on the location, distribution, and movement of EAB. Surveys also function as a delimiting tool to determine scope and extent of current EAB infestations. Survey information provides the basis for management decisions and provides continuous assessment of the effectiveness of the quarantine and control activities.

Trace Back and Trace Forward Investigations Surveys are conducted to find new infestations and determine the extent of a known infestation. In the course of conducting surveys or carrying out regulatory activities, investigations of the movement of potentially infested articles can also be considered a survey activity. Trace back investigations are conducted to try to determine the source of an infestation and trace forward investigations help determine if further spread occurred due to movement of infested host material or means of conveyance.

Activities that may require investigations in the EAB program include the movement of nursery stock, logs, lumber, wood chips, pallets or firewood.

General Detection Surveys The purpose of a general detection survey is to determine if a pest exists in an area. Positive results indicate that a pest is present. Lack of a positive result is valuable for providing clues to dispersal, temporal, or spatial activity patterns of pests particularly when considered with positive results from similar areas or proximities.

Early detection of isolated infestations discovered during general detection surveys may provide additional opportunities to manage the infestation. The size of the infestation cannot be determined, however until a delimiting survey is conducted.

The current program survey method for *Agrilus planipennis* is the selection of locations for the placement of panel traps. Attractant-baited panel traps offer several advantages over previous detection methods including lower cost, uniformity of sampling unit, safety, fewer logistical problems, and more precision in sampling.

Destructive sampling may be used to recover EAB larvae from suspect infestations identified from visual surveys for purposes of dendrochronology study.

Attractant-baited panel traps have been determined to be the most effective and operationally efficient method of survey (Marshall et al. 2009, Francese et al. 2006). Aerial survey and remote sensing

have not proven to be an efficient method of survey for EAB at this time.

Activities for conducting general surveys include:

- Planning, prioritizing, and procuring equipment and supplies for survey activities
- Developing or adapting existing protocols to meet new or unusual site-specific program needs
- Assisting scientists with the development and evaluation of new or improved survey protocols
- Following procedures for reporting new infestations and prompt specimen identification
- Maintaining survey records and maps
- Reporting survey results to management officials in a timely fashion

Many EAB infestations have been found by the public. As such, a strong outreach program is crucial for survey to be utilized efficiently. For current information on survey protocol, please consult the USDA Survey Guidelines located at:

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/index.shtml

Visual Survey

Although visual survey is not recommended as a primary survey tool for early detection, it can be helpful in locating older (3-5 yr.) infestations. The EAB Program uses visual survey to rapidly assess new infestations. Visible symptoms are difficult to determine in early infestation stages with no visible symptoms in the first year of infestation. Visual survey may only detect trees that have been infested for three or more years. Early stages, which show few signs of infestation, will not be detected by visual survey alone. Panel traps are most effective in detecting earlier stage infestations (Marshall, 2009). On larger trees, symptoms may be present only in the upper canopy in the early stages of infestation. In special instances, tree climbers may be used for a more intense inspection than a visual survey would provide from the ground.

Trees in a state of decline or with a combination of other symptoms should be examined more closely with a tree climber or a destructive sampling technique. They should be marked and their location recorded with GPS coordinates. If no life stages, exit holes, or serpentine galleries under bark are found, the tree will be considered negative for data recording purposes.

Symptoms of EAB Infestations Symptoms of *Agrilus planipennis* infestation to be cognizant of when conducting visual surveys include:

- Canopy stress/dieback



- Epicormic shoots/suckering



- Bark splits with larval galleries underneath



Figure 5.1 Symptoms of EAB on *Fraxinus* spp.

- Woodpecker Damage



- D-shaped exit holes (3-4 mm diameter)



- Serpentine larval galleries



Figure 5.2 Symptoms of EAB on *Fraxinus spp.*

Delimiting Survey

The purpose of a delimiting survey is to gather population density and dispersal information that will assist in planning a strategy for management.

- 1) After detecting EAB(s) in traps or finding an infested tree, conduct a visual survey until infested trees are no longer found. Continue visual survey for a distance of two miles beyond the initial trap catch or infested tree detection.
- 2) Determine the age of the infestation by examining the most heavily infested trees in the area (protocol to be provided by APHIS PPQ).
- 3) When phenologically appropriate, establish a systematic grid of traps starting at the last known infested trees and extending for one mile in radius for every year that the infestation was determined to have existed and then add one additional mile.

Example: Aging of the infestation indicates that it is three years old. From the last infested tree determined from visual survey, draw a band four miles wide (3 years + 1 mile) around the infestation and survey the area with program traps.

- 4) It is recommended trap deployment should be conducted with eight traps placed per square mile where ash trees are accessible.
- 5) If a candidate area is devoid of ash, the area should be omitted. Exceptions to setting traps in only ash trees include areas where volumes of potentially infested logs and/or firewood were introduced to the site.

If additional personnel are to be used, a training session for all participants should be organized. Information covered should include recognition of EAB life stages, damage symptoms, ash tree identification, description of infested site and survey methods, data collection protocols, and safety considerations.

Destructive Sampling

Destructive sampling of standing trees should be limited to specific trees displaying two or more symptoms of EAB in an uninfested county.

For optimal detection capability, special attention should be given when destructive sampling is done in the late summer and early fall months, as the first larval instars are especially difficult to detect due to their extremely small size.

Monitoring Survey The purpose of a monitoring survey is to evaluate the effectiveness of an action to contain or suppress EAB. Use the same survey tools used for delimiting surveys.

Survey Sample Collection Please refer to instructions provided in Section 3. Identification.

Quality Control for Survey Activities Survey activities will be subjected to quality control monitoring. Quality control techniques will include resurvey of a percentage of the sites, analysis of survey data, and by direct observation. Other methods to assure sensitivity and selectivity as well as the quality of the techniques employed should be developed.

Instructions To Officers

Officers must monitor and certify regulatory treatments or other procedures when authorizing the movement of regulated articles. Only authorized treatments may be used in accordance with agency manuals and/or labeling restrictions.

Authorities

The Plant Protection Act of 2000 (Statute 7 USC 7701-7758) provides authority for implementation of emergency quarantine action. This provision is for interstate regulatory action only; intrastate regulatory action is provided under state authority. State departments of agriculture normally work in conjunction with federal actions by issuing their own parallel hold orders and quarantines for intrastate movement.

Emerald ash borer is not considered an extraordinary emergency as defined by the Plant Protection Act. However, if the U.S. Secretary of Agriculture determines that an extraordinary emergency exists and that the measures taken by the state are inadequate, USDA can take intrastate regulatory action provided that the governor of the state has been consulted and a notice has been published in the *Federal Register*. If intrastate action cannot or will not be taken by a state, then PPQ may find it necessary to quarantine an entire state.

The Federal Emerald Ash Borer (EAB) Quarantine (7CFR 301.53) provides the authority to conduct regulatory activities. In addition to the Federal Quarantine, individual states have established state quarantines regulating intrastate movement of regulated material in accordance with federal regulations.

PPQ works in conjunction with states' department of agriculture to conduct surveys, enforce regulations, and take control actions. PPQ employees must have permission of the property owner before entering private property. Under certain situations during a declared extraordinary emergency or if a warrant is obtained, PPQ can enter private property in the absence of owner permission. PPQ prefers to work with the state to facilitate land access. However, each state government has varying authorities regarding entering private property. A general Memorandum of Understanding (MOU) developed between PPQ and each cooperating state refers to how cooperative program activities are conducted. For clarification, check with your USDA Program Manager, State Plant Health Director (SPHD) or State Plant Regulatory Official (SPRO) in the affected state.

Regulating the Spread of EAB

The rate of natural dispersal by adult flight of *Agilus planipennis* is estimated to be as little as 800 meters per year (Taylor et al, 2004). However, unpublished data suggests that this rate may be much higher and variable due to environmental factors and insect population density.

Tethered flight in laboratory conditions predicts that a mated female may fly more than 20 km (Taylor et al. 2006). Regulatory measures are necessary to prevent human-assisted spread through the transportation of *Agrilus planipennis* in firewood, logs, timber and other wood materials. Quarantine regulations typically prohibit the movement of potentially infested host material out of the infested area or require treatments, inspections, or other mitigating measures.

Regulated Articles

Regulated articles for *Agrilus planipennis* include:

1. The emerald ash borer (all life stages)
2. Firewood: all non-coniferous species
3. Nursery stock, green lumber, and other material living, dead, cut or fallen including logs, stumps, roots, branches, and composted and uncomposted chips of the genus *Fraxinus*.
4. Any article, product, or means of conveyance not listed above if an inspector determines that it presents a risk of spreading emerald ash borer and notifies the person in possession of the article, product, or means of conveyance that it is subject to the restrictions of the regulations.

Approved Regulatory Treatments

Approved regulatory treatments are determined by program management and/or a Technical Advisory Committee.

Table 6.1. Interstate movement of regulated articles from quarantined areas.

Regulated Article	Mitigation and Treatment Measures
Firewood of all hardwood species	<ul style="list-style-type: none"> • Remove bark and an additional ½ inch of wood¹ or • Kiln sterilization treatment (T404-b-4) or • Heat treatment (T314-a) in a heat treatment facility approved by APHIS or • Fumigate according to treatment schedule T404-b-1-1 (Methyl bromide fumigation at NAP-tarpaulin or chamber) or • Apply an APHIS approved method².
Chips and Mulch of all hardwood species	<ul style="list-style-type: none"> • Chip or mulch to less than one inch in at least two dimensions or • Follow an APHIS approved mulching or composting protocol or • Apply an APHIS approved method².
Nursery Stock of <i>Fraxinus</i> spp. Itself	No treatment available. No compliance agreements. No certification. This article is not being moved at this time.
Green lumber of <i>Fraxinus</i> spp. itself	<ul style="list-style-type: none"> • Remove bark and an additional ½ inch of wood¹ or • Kiln sterilization treatment (T404-b-4) or • Fumigate according to treatment schedule (404-b-1-1 (Methyl bromide fumigation at NAP-tarpaulin or chamber) or

6. Regulatory Procedures

Emerald Ash Borer

	<ul style="list-style-type: none"> • Apply an APHIS approved method².
Logs of <i>Fraxinus</i> spp. itself	<ul style="list-style-type: none"> • Remove bark and an additional ½ inch of wood¹ or • Kiln drying treatment (T404-b-4) or • Heat treatment (T314-a) in a heat treatment facility approved by APHIS or • Fumigate according to treatment schedule T404-b-1-1 (Methyl bromide fumigation at NAP-tarpaulin or chamber) or • Apply an APHIS approved method².
Other material including wood waste, living, dead, cut or falling including stumps, roots, branches of <i>Fraxinus</i> spp.	<ul style="list-style-type: none"> • Chip or mulch to less than one inch in at least two dimensions or • Apply an APHIS approved method².
WPM containing regulated green lumber, including but not limited to , dunnage, crating, pallets, packing blocks, drums, cases, and skids.	<ul style="list-style-type: none"> • ISPM accredited treatments or • Treatment/mitigations for green lumber of <i>Fraxinus</i> spp. itself as listed above or • Apply an APHIS approved method².
<p>¹ The bark and wood removed will be regulated separately. If intended for interstate movement the removed bark and wood must be treated as described in Table 1 for chips and mulch. If produced at a mill located outside the quarantine area but approved to handle green ash logs or lumber from within the quarantine area, wood waste must be treated or destroyed prior to adult flight season.</p> <p>² Consult Local USDA APHIS Official or State Plant Health Director.</p>	

Detailed specifications for treatments can be found in the current edition of the PPQ Treatment Manual online at:

http://www.aphis.usda.gov/import_export/plants/manuals/ports/treatment.shtml

Quarantine Boundaries

Quarantine boundaries will be determined in consultation with state authorities based on the following factors:

- Human-assisted (passive) dispersal
- Natural (active) dispersal
- Ease of enforcement due to geopolitical boundaries
- Historical and existing quarantine practices

Regulatory Management of Outlying Infestations

Isolated infestations outside of the quarantined boundary are common occurrences in pest management programs. These isolated infestations generally represent anthropogenic (human-assisted) dispersal of *Agrilus planipennis* created by movement of infested articles.

Regulatory management of outlying infestations will be handled using the following procedures:

1. Upon confirmation by a USDA EAB identifier and with consensus from the cooperating regulatory agencies, a quarantine will immediately be established. The quarantined area may be adjusted based on additional survey information.
2. A delimiting survey will be initiated as soon as possible to establish the area of impact.
3. Provisions of the temporary quarantine will be the same as those established in the formal quarantine. Newspapers and direct mailings will be used to notify inhabitants within the established area that a temporary quarantine is in effect.
4. If the cause of the outlying infestation is not readily determined, trace back inspections and interviews with local business and home owners will be conducted to determine the source. These inspections will begin at the epicenter of the infestation and work outward. Trace forward inspections will be conducted to determine if infested host material has been moved out of the area. These inspections should include all establishments located in and/or conducting business within the regulated area.

Regulated Establishments

Field personnel will attempt to detect the pest within the regulated area at all establishments where regulated articles are sold, grown, handled, moved, or processed. Involved establishments might include, but are not limited to:

- Campgrounds
- Firewood dealers
- Nurseries
- Logging/lumber companies
- Pallet manufacturing companies
- Landscapers and garden centers
- Utilities companies
- Tree removal companies
- Municipality forestry services
- Waste removal companies

Please refer to Appendix F for additional sources to identify regulatory concerns.

Principle Regulatory Activities

Regulatory activities are directed at implementing and enforcing quarantine provisions governing movement of regulated articles which could result in human-assisted dispersal of *Agrilus planipennis*. Typical activities include:

1. Identifying persons and establishments whose business or personal activities could result in the further dispersal of *Agrilus planipennis*. (See Regulated Establishments, above, and

- Appendix F.)
2. Contacting by regulatory visit, mail, email, and/or telephone those identified persons or establishments to explain quarantine provisions.
 3. Determining if provisions of the quarantine (e.g., treatment or processing) may be applied to permit the person or establishment to move regulated articles out of the quarantined area.
 4. Conducting inspections of regulated articles and monitoring procedures to mitigate pest risk on a shipment by shipment basis.
 5. Issuing Limited Permits (PPQ form 530) to allow movement of regulated articles out of the quarantine area to a specific destination for further processing or treatment.
 6. Issuing Certificates (PPQ form 540) to allow movement of regulated articles out of the quarantined area when they have been treated or processed in such a manner that they no longer present a risk for further dispersal of *Agrilus planipennis*.
 7. Entering into a Compliance Agreement at the discretion of the inspector with a person or establishment that can demonstrate their ability to meet the provisions of the quarantine.
 8. Conducting periodic physical site visits to monitor the activities of those persons or establishments placed under a Compliance Agreement to observe and assess treatments or other processes and activities. Periodicity of monitoring is based on risk assessment of the product and the business or individual under compliance.
 9. Conducting and/or assisting with investigations of suspected violations of the quarantine as necessary and appropriate.
 10. Recording information about contacts, visits, and Compliance Agreements and maintaining data on persons or establishments affected by the quarantine.
 11. Reporting results of regulatory activities to management officials on a weekly basis.
 12. Conducting special regulatory operations.
 13. Researching local and interstate trade movement to determine pathway risk.

Investigations and Violations

When regulated material is suspected to have been moved out of the regulated area in violation of the quarantine, regulatory personnel will conduct initial preliminary investigations to determine if a violation of the quarantine has occurred and safeguard any regulated material. These investigations will also attempt to identify and to trace the source and destination of any other related shipments of regulated materials that have occurred.

Preliminary investigations by regulatory personnel will allow management to determine whether the situation warrants additional

formal investigation by USDA-APHIS-Investigation and Enforcement Services (IES) personnel.

Regulatory Records

All data related to regulatory activities will be maintained in the regulatory database. Regulatory personnel will record information such as:

1. Date of visit
2. Purpose of visit (routine, investigation, monitoring) and actions taken/needed
3. GPS coordinates of the site
4. Name and address of contact
5. Phone, fax, and/or e-mail of contact
6. Communication method
7. Type of contact
8. Compliance agreement number (if applicable)

Weekly status reports will be made by regulatory personnel to EAB management officials summarizing their activities and highlighting immediate and developing problems.

Quality Control

Management officials will review the results of visits, contacts, and Compliance Agreements maintained by regulatory personnel to ensure that program standards are followed. A percentage of establishments operating with or without a Compliance Agreement will be selected for a paper audit on a regular basis. This is important to reinforce the effectiveness of a Compliance Agreement.

Outreach

Outreach is a vital component of every aspect of the EAB program. Without public support and cooperation, the efficacy of the program is very limited. Regulatory personnel should utilize opportunities during general regulatory activities and special regulatory operations to inform the public about the EAB program and enlist their cooperation.

Special Regulatory Operations

Another aspect of regulatory activity is to identify and conduct special operations to serve as deterrents and quality control for movement of regulated articles. These operations also give unique opportunities to inform the public of the pest and related regulations. The operations may be conducted with state cooperators. Special regulatory operations may include:

- Highway operations in cooperation with local law enforcement which stop traffic to inspect for movement of regulated articles by private citizens and commercial dealers and to increase public awareness.
- Rest stop and weigh station operations to monitor quality

control.

- Ride along operations with law enforcement to regulate and ensure commercial enterprise compliance.
- Campground operations to regulate the movement of firewood from quarantined areas.
- Ferry operations to monitor the movement of regulated articles.
- Special focus operations to concentrate regulatory activities in a smaller area of interest.

Any special regulatory operation should be coordinated with tribal governments and federal, state, and local authorities including, but not limited to: tribal, state, and local police departments, states' department of agriculture, states' department of transportation, states' department of natural resources, and local wood disposal and utilization entities.

Issuing an Emergency Action Notification

After an initial suspect positive detection in a new area outside a quarantined area, an Emergency Action Notification (PPQ form 523) may be issued to place a hold on regulated material or facilities where regulated material may be found pending positive identification by a USDA-APHIS-PPQ recognized authority.

An Emergency Action Notification (EAN) may also be issued within the quarantine area to order a hold, treatment, destruction or other safeguarding action for a regulated material due to a potential pest risk. The EAN may be completed by hand in the field, but must be transferred into the EAN database within 14 days. Authority for EAN issuance is to be determined by each state's command structure.

Removing areas from Quarantine

Project managers identify and remove areas from quarantine requirements after *A. planipennis* is declared eradicated. Quarantined areas will be released following three years of negative survey after eradication is declared. Release of the quarantined area will occur with the consensus of the cooperating regulatory agencies. APHIS will publish a Notice of Quarantine Revocation in the Federal Register when areas are removed from quarantine requirements.

Regulatory Forms

Sample compliance agreement templates can be found at the following link:

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/regulatory.shtml

Other sample regulatory forms and useful documents are located in the appendices and include:

- Info Packets

- Regulatory operation forms
- Sources to find regulatory concerns
- PPQ 518 form (Report of Violation)
- PPQ 530 form (Limited Permit)
- PPQ 540 form (Certificate)
- PPQ 523 form (Emergency Action Notification)

Overview

The Emerald Ash Borer Program has transitioned from an eradication program to a management program. Effective and cost efficient control technologies are not currently available to apply area-wide to effect pest eradication. In the future additional tools may become available to suppress the dispersal of the pest. Program partners are conducting extensive research to develop additional tools and methodologies.

This section provides the program policy on management of isolated infestations and currently available information on control and management tactics. This will assist states, local communities, and the public in mitigating the effects of this destructive pest.

As research is ongoing in this area, management procedures will be subject to frequent changes. Please contact EAB Program Management to ensure that procedures are up to date before implementing management measures. Further guidance may be found in the APHIS Emergency Programs Manual at:

http://www.aphis.usda.gov/import_export/plants/manuals/emergency/downloads/epm.pdf

Community Preparedness Planning

Since control options to eradicate or prevent dispersal are not effective at this time, it is recommended that communities take action to prepare for an infestation of emerald ash borer.

Examples of preparedness/response plans for states may be found at:

<http://www.emeraldashborer.info/communityplan.cfm>

Examples of individual city response plans may be found at the following URLs:

<http://www.dnr.state.oh.us/Portals/18/eab/pdf/EAB%20plan%20Marietta.pdf>

<http://dnr.wi.gov/forestry/uf/eab/EABToolkitCD/03.%20Planning/e.%20Community%20Examples/EAB%20Management%20Plan%20-%20Upper%20Arlington%20OH.pdf>

<http://www.agr.state.il.us/eab/data/200807303570.pdf>

Forest management experts suggest taking preventive measures such as diversifying landscape plantings using the 30:20:10 rule: The urban forest should be composed of no more than 30% of the same family, 20% of the same genus, and 10% of the same species. Choosing native

species over non-native species is recommended in order to reduce costs associated with maintenance (watering and fertilizers). Ash (*Fraxinus* spp.) should be avoided as landscape plantings near infested areas/states.

Models exist for ash phloem reduction in managed forests/woodlots. One such model can be found at: <http://www.ashmodel.org/> Reduction of ash populations should be considered in consultation with forest management experts.

It is prudent to have a response plan prior to an EAB infestation. An example of a response planning exercise may be found at:

<http://www.mda.state.mn.us/news/publications/pestsplants/insects&pests/eab/eabxrciserpt.pdf>

Biological Control

In order to be a successful biological control agent, the following traits are desired:

1. Parasitoid (organism that lives on or in another organism)
2. Multivoltine (producing more than one generation per year)
3. Monophagous (feeding on only one type of food)

Three parasitoids that meet the above criteria are currently under evaluation by APHIS. These parasitoids are known to attack EAB consistently in its native range in China. The biocontrol agents include one larval ectoparasitoid, *Spathius agrili* (Hymenoptera: Braconidae) (Yang et al. 2005), one species of egg parasitoid, *Oobius agrili* (Hymenoptera: Encyrtidae) (Zhang et al. 2005), and one species of larval endoparasitoid, *Tetrastichus planipennisi* (Hymenoptera: Eulophidae) (Liu et al. 2003; Yang et al. 2006).

Research on EAB biological control began in 2002 when this destructive beetle was first found in Michigan. Much of the biocontrol research was done in China, where studies of EAB led to the discovery of three specific natural enemies: *Spathius agrili*, *Oobius agrili* and *Tetrastichus planipennisi*. These natural enemies are tiny stingless wasps that seek and kill EAB eggs and larvae. Five years of research led to an environmental assessment of field release of these natural enemies. After a 60-day public comment period and a “Finding of No Significant Impact”, APHIS and the State of Michigan approved release of these wasps for control of EAB. Small-scale inoculative releases were performed in Michigan in 2007 and 2008. The sites will continue to be monitored for establishment of the natural enemies and evaluated for their potential to control and slow the dispersal of EAB in the United States. Early results indicate that two species are capable of reproducing in North America.

An EAB biocontrol laboratory has been established at the APHIS facility in Brighton, Michigan. Tools and techniques for rearing natural enemies will be further developed at this location with the goal of future distribution of parasitoids to other states. Over the next five years, specific release sites will be designated for research and tracked to collect data to determine:

- successful establishment of natural enemies
- numbers of natural enemies needed for establishment
- site conditions required for establishing natural enemy populations
- interactions among the three exotic natural enemies and native natural enemies
- natural enemy dispersal rates
- impacts on EAB populations and ash survival or recovery
- effects on non-target species

At the conclusion of the first five years of the EAB biological control program, scientists will evaluate which natural enemies are most suitable for long-term mass production and area-wide release.

With oversight from APHIS and the U.S. Forest Service, the rearing facility will be responsible for rearing these three natural enemies for scheduled releases. Release sites will be determined by APHIS and the U.S. Forest Service in consultation with state partners. The releases will depend upon program needs, regulatory approval, and production capabilities. The challenges of rearing these natural enemies will limit their availability for release until the laboratory is fully operational.

Contact: Ken Bloem, Biological Control Coordinator, 919-855-7407
email: Kenneth.Bloem@aphis.usda.gov

Initial releases and post-release monitoring, including impacts on EAB and non-target wood-boring beetles, and dispersal and establishment of each parasitoid species, will be conducted. The Environmental Assessment for this proposed release can be found at the following web site:

http://www.aphis.usda.gov/plant_health/ea/downloads/eab-ea4-07.pdf

Chemical Control

At this time available chemical treatments are not cost effective for large scale implementation. Therefore, the EAB program can not recommend chemical control on a large scale. However, depending on beetle population densities, recent research suggests individual trees may be effectively treated (Cappaert et al. 2006). Consult your county or university extension agent for information on approved treatments for your specific area.

If treatments selected or proposed are not in conformance with current pesticide labels, an emergency exemption can be requested and obtained under Section 18, or 24(c), special local need (SLN), of FIFRA, as amended.

**Eradication
Criteria**

Eradication Criteria Guidelines outside the current leading edge:

All new detections outside the leading edge will be evaluated individually and any management action supported by APHIS will be considered only after all the following criteria are met:

1. The outlier must be a single, clearly identifiable regulatory incident. This may include nursery stock, firewood, or other regulated material transported beyond the current buffer.
2. The population, using dendrochronology and a delimiting survey, must be demonstrated to be less than two years old or to have no more than one population release from the original host material.
3. The population, using current technologies and methodologies, must be identified as eradicable by the EAB Management Team (i.e., the point of introduction clearly identified and age of infestation as delineated in item 2).

The EAB Management Team may consider control actions for EAB populations older than two years if circumstances indicate eradication may be achieved or the control action is determined to be strategically important for slowing the spread.

**Management
Records**

Program personnel will maintain records and maps noting the locations of all detections, the number and type of management treatments, and the materials used in each treated area.

Monitoring

Program personnel will implement an effective monitoring program to evaluate program efforts, pesticide use, and environmental impact. The monitoring program also will address any concerns of APHIS, cooperators, and the public while providing data for assessment of the application method.

The monitoring program must be a combined effort between the state in which the emergency program is being conducted and PPQ. Specific tasks must be coordinated for monitoring activities.

- Overview** Environmental Services (ES) is a unit of APHIS' Policy and Program Development Staff (PPD). ES manages the preparation of environmental documentation, such as environmental impact statements and environmental assessments, to aid in program operational decisions. ES also coordinates pesticide registration and approvals for APHIS pest control and eradication programs, ensuring that registrations and approvals meet program use needs and conform to pesticide use requirements. Contacts and informational websites are listed in the Resources Section of this document.
- Disclaimer** All uses of pesticides must be registered or approved by appropriate Federal, State, and/or Tribal agencies before they can be applied. The information provided on pesticide labels may not reflect all of the actual information, including precautions and instructions for use, which you are required to follow in your specific State or locality. It is the responsibility of persons intending to use a pesticide to read and abide by the label, including labeling that has been approved for the particular State or locality in which the chemical is to be used, and to comply with all Federal, State, Tribal, and local laws and regulations relating to the use of the pesticide. APHIS program staffs are responsible for their compliance with applicable environmental regulations.
- Protected Species** To ensure that protected species are not impacted negatively during survey and control activities, contact the Emerald Ash Borer (EAB) Program environmental monitor.
- To date, program activities have significantly interfaced with several protected species. Specific protocols have been implemented to mitigate negative impacts on the Indiana gray bat, the bald eagle, the copperbelly water snake, and other species in consultation with Fish and Wildlife Services.
- For further details, see the biological assessment and contact the EAB program environmental monitor.
- Environmental Assessment** All program control activities require review by Headquarters Environmental Services staff before any work may begin. Control activities include eradication and containment actions that require cutting and/or destruction of trees. An environmental assessment is mandatory before APHIS supported control or containment activities may begin.

Introduction

Education and outreach plays a key role in communicating the operational objectives of the EAB Program; detection, control and eradication. Informed and supportive targeted audiences (industry stakeholders, congressional constituents, special interest groups and the public in general) can serve as advocates as well as active participants in the Cooperative Emerald Ash Borer Program.

It is never too early to begin the education and outreach process. To facilitate the process, the EAB Program has many generic outreach materials that are available and free of charge. **To prevent duplication of efforts and to ensure consistency, contact the EAB program's Legislative and Public Affairs (LPA) specialist to review existing outreach materials. Furthermore all new communication tools (publications, videos, posters, news releases, etc) where the USDA is mentioned, must be reviewed by the LPA program specialists prior to publication.**

General Outreach Objectives

1. Coordinate outreach efforts among cooperators to ensure the program has a consistent message.
 - A. Establish a small, core committee, to manage and direct program communication; membership should include local, state, and federal partners, cooperative extension and other partners deemed relevant.
 - Identify roles and responsibility for individuals and as a whole
 - Identify policies and procedures for releasing information to public and stakeholders
 - Address financial responsibilities
 - Develop a two-way reporting mechanism with Cooperative Management Staff
 - Establish a Communications Advisory Committee and identify roles and responsibilities. Membership could include industry groups such as professional landscapers and arborists, lumber and wood industry, nursery owners, environmental organizers, and other economically impacted groups. This group will be an invaluable network for the dissemination of messages and information.
2. Develop outreach materials to meet various program needs and reach various audiences. Outreach personnel will work with EAB program managers to identify areas or activities where outreach materials are needed and do not currently exist.

Priorities will be as follows:

- Identify target audiences and leadership
 - Identify key messages for all program initiatives; survey, public meetings, tree removal and other control activities, etc.
 - Identify key messages to support regulatory activities: “Don’t move firewood”, examine your trees, know federal and state regulations, etc.
3. Deliver outreach materials through a variety of outlets to ensure widest exposure.
- Create a marketing mix using mainstream/alternative/grassroots media to reach targeted audience. (Television, radio, outdoor, newspaper, internet, industry publications, civic groups, newsletters, etc.)
 - Reach out to specialized target audiences through their industry newsletters, web site links, and association meetings. Develop “champions” for the program.
 - Seek out opportunities for communication: country fairs, home and garden shows, regional association meetings, Chamber of Commerce events, etc.
 - Establish a speakers’ bureau and create canned power point presentations.

General Outreach Activities and Initiatives

Some of the common activities that might be expected in the outreach portion of an EAB program include:

1. Maintain the EAB website to provide access to current information on EAB, quarantines, survey areas, etc.
2. Maintain a toll-free EAB hotline.
3. Catalogue and review existing informational materials to prevent duplication of effort and ensure consistency.
4. Develop public service announcements and arrange for broadcast on radio and/or television. Utilize a spokesperson with name recognition, if possible.
5. Keep the media informed regarding program activities.
6. Keep local officials, local government, community leaders informed about the program
7. Craft easy to read letters regarding time, date, location, and purpose of public meetings and execute mailing
 - Use tax rolls for names and addresses (Note: Tax rolls apply to owners not necessarily residents.)
 - Allow substantial lead time.
 - In resort areas, allow for absentee homeowner issues.
8. Arrange, moderate, and provide presentations and support at public meetings

9. Periodically meet with program staff for program feedback, problems, concerns, etc.
10. Create specialized communication vehicles when needed (magnets, tattoos, stickers, etc.) to support EAB awareness
11. Continually refine and develop communication vehicles (brochures, posters, newsletters, etc.) to ensure accuracy and current program information.
12. Develop and arrange for publication of news releases for mainstream and alternative media
13. Craft easy-to-read letters regarding invasive program activities (survey, tree marking, eradication, etc.) and execute mailings.

Outreach Material There are generic program materials to assist in public education. To prevent duplication of effort and to ensure consistency, before implementing outreach activities contact the EAB LPA specialist to review existing outreach materials.

Telephone Hotline The toll-free telephone National EAB hotline is 1-866-322-4512. The hotline is staffed by trained and knowledgeable personnel who can answer questions about the EAB program and direct callers to appropriate personnel for more in depth questions. Statistical information, such as the number and nature of calls, should be reported to outreach personnel.

Web Site www.emeraldashborer.info is the national website developed by the Cooperative EAB Program and funded by the U.S. Forest Service as a source and link to federal and state information. It contains current program information including: quarantine information, maps, EAB signs and symptoms, ash tree identification, treatment options, prevention, tree removal and replacement and on-going research.

Additionally, general public may email the program at EmeraldashborerInfo@aphis.usda.gov.

Public Meetings or Informational Open Houses Public meetings or informational open houses will take place when deemed necessary and appropriate in particular when control efforts such as tree removal or pesticide treatments impact a community. These meetings will address public concerns, communicate the program strategy and actions, and secure community support.

Outreach personnel will coordinate scheduling and facilities, ensure that public meetings or informational open houses notifications are posted in appropriate newspapers and other media outlets and that handouts, fact sheets, informational posters, and other outreach materials are readily available at the meeting. Where possible, outreach personnel will make advance direct mailings to notify residents of the impacted community

of the meeting.

Public meetings will be organized as deemed appropriate and will include partner agency participation such as:

1. Political representatives and community leaders who are familiar with local concerns and recognized by the local community.
2. State and Federal program representatives who can respond to questions about EAB, quarantine restrictions, control measures, and its impact.
3. Representatives from cooperating state universities who can answer questions about biology of EAB, its host range, and potential impact in the United States.
4. County, city and local cooperators who can respond to questions about their roles.

Media Relations

The APHIS PPQ and LPA staff should be notified as soon as possible after EAB is confirmed and routinely notified of any media requests. All national media calls must be coordinated with APHIS LPA. One primary media spokesperson should be designated by the cooperator to work with the EAB LPA spokesperson. To avoid conflicting and confusing statements, all outgoing information should be processed through the designated spokesperson. The spokesperson should thoroughly understand particular aspects of the program, such as survey, regulatory, and control activities, as well as the federal and state quarantines. All EAB program spokespersons will develop and maintain close contacts with each other and reporters and community group leaders to provide accurate and consistent information. If no personnel at the local level exist or he/she does not have adequate media experience, the EAB LPA specialist should be notified so he/she can provide experienced media representation to the program.

Literature cited:

- Akiyama K. and S. Ohmomo. 1997. A checklist of the Japanese Buprestidae. Gekkan-Mushi (Supplement 1).
- Bauer, L. S., R. A. Haack, D. L. Miller, T. R. Petrice, and H. Liu. 2004. Emerald ash borer life cycle. In: *Emerald ash borer research and technology development meeting*, Compiled by: V. Mastro and R. Reardon.
- Cappaert, D., D. G. McCullough, T. M. Poland, and N.W. Siegert. 2005. Emerald ash borer in North America: A research and regulatory challenge. *American Entomologist*, 51:152-165.
- Chinese Academy of Science, Institute of Zoology. 1986. *Agrilus marcopoli* Obenberger. Agriculture Insects of China (part I), China Agriculture Press, Beijing, China.
- Francese, J. A., I. Fraser, D. R. Lance, and V. C. Mastro. 2006. Developing survey techniques for emerald ash borer: The role of trap height and design. In: *Emerald ash borer and Asian long-horned beetle research and technology development meeting*, Compiled by: V. Mastro, D. Lance, R. Reardon, and G. Parra.
- Liu, H-P, L.S. Bauer, R-T Gao, T-H Zhao, T.R. Petrice, and R.A. Haack. 2003. Exploratory survey for the emerald ash borer, *Agrilus planipennis* (Coleoptera: Buprestidae), and its natural enemies in China. *Great Lakes Entomologist*. 36:191-204.
- Lyons, D. B. G. C. Jones, and K. Wainio-Keizer. 2004. The biology and phenology of the emerald ash borer. In: *Emerald ash borer research and technology development meeting*, Compiled by: V. Mastro and R. Reardon.
- MacFarlane, D. W. and S. P. Meyer. 2005. Characteristics and distribution of potential ash tree hosts for emerald ash borer. *Forest Ecology and Management*, 213:15-24.
- Marshall, J. M., A. J. Storer, I. Fraser, and V. C. Mastro. 2009. Efficacy of trap and lure types for detection of *Agrilus planipennis* (Col., Buprestidae) at low density. *Journal of Applied Entomology*, In press.
- McCullough, D. M. and N. W. Siegert. 2006. Using girdled trap trees effectively for emerald ash borer detection, delimitation, and survey. <http://www.emeraldashborer.info/files/handoutforpdf.pdf> Last accessed January 18, 2008.
- McPartlan, D., P. D. Bell, C. Kellogg. 2006. Eradication of emerald ash borer in

- Michigan, Ohio, and Indiana – Implementation of the strategic plan. January 11, 2006 revision. http://www.aphis.usda.gov/plant_health/ea/downloads/eab-ea4-07.pdf Last accessed January 18, 2008.
- Nomura, S. 2002. *Agrilus planipennis*. Canadian Food Inspection Agency Science Branch. Last accessed November 30, 2004.
- Poland T. M. and D. G. McCullough. 2006. Emerald ash borer: Invasion of the urban forest and the threat to North America's ash resource. *Journal of Forestry*, 118-124.
- Sugiura N. 1999. The family Buprestidae in Fukushima Prefecture: the genus *Agrilus*. On-line at: <http://www1.linkclub.or.jp/~sugin/fukusima/nagatama/nagatama2.html> Last accessed August 2002.
- Taylor, R. A. J., T. M. Poland,, L. S. Bauer, K. N. Windell, and J. L. Kautz. 2006. Emerald ash borer flight estimates revised. In: *Emerald ash borer research and technology development meeting*, Compiled by: V. Mastro, D. Lance, R. Reardon, and G. Parra.
- Taylor, R. A. J., L. S. Bauer, D. L. Miller, and R. A. Haack. 2004. Emerald ash borer flight potential. In: *Emerald ash borer research and technology development meeting*, Ed. V. Mastro and R. Reardon.
- Timms, L. L., S. M. Smith, and P. de Groot. 2006. Patterns in the within-tree distribution of the emerald ash borer *Agrilus planipennis* (Fairmaire) in young green-ash plantations of south-western Ontario, Canada. *Agricultural and Forest Entomology*, 8:313-321.
- US Forest Service. 2008. Emerald ash borer maps. <http://www.ncrs.fs.fed.us/4501/eab/maps/> Last accessed: 01/17/08.
- Yang, Z-Q., J.S. Strazanac, Y-X Yao, and X-Y Wang. 2006. A new species of emerald ash borer parasitoid from China belonging to the genus *Tetrastichus* Haliday (Hymenoptera: Eulophidae) parasitizing emerald ash borer from China. *Proceedings of the Entomological Society of America*. 108:550-558.
- Yang, Z-Q, J. S. Strazanac, P. M. Marsh, C. van Achterberg, and W-Y, Choi. 2005. First recorded parasitoid from China of *Agrilus planipennis*: A new species of *Spathius* (Hymenoptera: Braconidae, Doryctinae). *Annals of the Entomological Society of America*. 98(5):636-642.
- Yu, C. 1992. *Agrilus marcopoli* Obenberger (Coleoptera: Buprestidae), pp. 400-401. In:

Forest Insects of China (2nd edition) Ed. G. Xiao, China Forestry Publishing House, Beijing, China.

Zhang, Y-Z., D-W Huang, T-H Zhao, H-P Liu, and L.S. Bauer. 2005. Two new egg parasitoids (Hymenoptera: Encyrtidae) of economic importance from China. *Phytoparasitica*. 33:253-260.

Anthropogenic	Human-caused
APHIS	Animal and Plant Health Inspection Service
Cambium	The meristematic tissue in woody plants that exists between the wood (xylem) and the inner most bark (phloem)
CFR	Code of Federal Regulations
CPHST	Center for Plant Health Science and Technology
Cuneiform	Triangular or wedge shaped
DBH	Diameter at breast height
Delimiting survey	A sampling method to determine extent of an infestation of an exotic species in an area
Destructive sampling	Method of observing signs and symptoms of the presence or absence of a pest by destruction of the living sample unit; <i>i.e.</i> , removal of bark to look for larvae
Detection survey	A sampling technique to determine the presence or absence of a non-native species in an area
Detection tree technique	Determining the presence or absence of a pest through the use of randomly or strategically placed host trees artificially manipulated to attempt attraction of target pest organisms. These trees may be maintained and serviced on a schedule dictated by the goal of the survey
Developmental thresholds	The minimum and/or maximum temperatures that support physiological development of a species
Diapause, Facultative	The cessation of growth and reduction of metabolic activity in a species which occurs seasonally or when environmental conditions are unfavorable
Diapause, Obligatory	A period of quiescence genetically controlled and affecting every individual of every generation within a species regardless of environmental conditions
Dispersal, Active	The spread of an organism by its own method of locomotion (<i>e.g.</i> , walking, flight, etc.)
Dispersal, Passive	The spread of an organism aided by other than its own method of locomotion (<i>e.g.</i> , wind, water, man, etc.)

EAB	emerald ash borer
EAN	Emergency Action Notification
Ectoparasitoid	A parasitoid that develops outside the host and is attached or embedded in host tissue
Endoparasitoid	A parasitoid that develops inside the host
Epicormic shoots	Fast growing, relatively soft stems arising from the main trunk below the crown or root collar of a dying or stressed tree. Sometimes referred to as “water sprouts” or “suckering”
ER	Eastern Region
ES	Environmental Services
Exotic species	An organism or pest species not native to nor historically resident in North America, also referred to as alien, non-native, or invasive species
F	Fahrenheit
FIFRA	Federal Insecticide, Fungicide and Rodenticide Act
FY	Fiscal year
Generally infested area	All areas within a line extending from the peripheral positive EAB finds
Generation	One complete life cycle
GPS	Global positioning system
Growing degree day	A measure of the number of thermal units (degrees) that accumulate above a specified base temperature over a 24 hour period
Host	A species that provides food, shelter, or reproductive requirements for another organism
IES	Investigative and Enforcement Services
ISIS	Integrated survey information system
ISPM	International Standard for Phytosaniatry Measures

km	kilometer
Leading edge	The boundary defined by the line delineating the generally infested area
LPA	Legislative and Public Affairs
mg	milligram
mm	millimeter
Monophagous	Feeding on only one type of food
MOU	Memorandum of understanding
Multivoltine	Producing more than one generation per year
NC	North Central Forest Service Station
Obscure-aeneous	Dark coppery color
Parasitoid	An organism that lives on or in another organism, usually referred to as its host, and from which it obtains nourishment
PDA	Personal digital assistant
Phloem	Nutrient conducting tissue of the inner bark
Phototactic	Movement of an organism toward or away from a light source
PPD	Policy and Program Development
PPQ	Plant Protection and Quarantine
Regulated articles	All known or suspected hosts of a confirmed infestation of a non-native species, including soil and any other suspected product or article
SEL	Systematics Entomology Laboratory
Semio-chemicals	Chemicals emitted by plants, animals and other organisms and synthetic analogues of such substances, that evoke a behavioral or physiological response in individuals of the same or other species
SLN	Special local needs
SPHD	State Plant Health Director

SPRO	State Plant Regulatory Official
Thanatosis	Feigning death in order to avoid predation
Thermotactic	Movement of an organism toward or away from a heat source
USDA	United States Department of Agriculture
USFS	United States Forest Service
Visual survey	Simple examination of areas for eggs, larvae, pupae, or other evidence to determine if a particular insect species is present
VPN	Virtual private network
WPM	Wood packing material
WR	Western Region
Xylem	Water conducting tissue that comprises the bulk of most woody plants; wood

Agrilus planipennis Fairmaire Screening Aid

J. E. Zablony, USDA APHIS PPQ



Figure 1. *Agrilus planipennis* Fairmaire.

Members of the genus *Agrilus* are challenging to identify due to structural coloration and subtle morphological differences between species. Furthermore, the presence of newly discovered exotic *Agrilus* species in the Midwest and Ontario complicates identification issues and demands a renewed interest in applied taxonomy of *Agrilus* beetles.

The emerald ash borer (EAB) is an invasive buprestid beetle native to Northeast China, Korea, Japan, and Russia (Figure 1). In Michigan, and Ontario, this pest is established and has devastated stands of native ash trees. The following characteristics can be used for discriminating EAB from other native *Agrilus* species:

- No patches of pubescence
- Pronotum copper/green
- Elytra and abdominal sternites emerald green

- Abdominal tergites purplish copper in color
- Length 13 mm
- Emarginate pygidial spine (Figure 2).
- **EXOTIC**



Figure 2. Pygidium of *A. planipennis*.




Figure 3. Aedeagus of *A. planipennis* (a. tergal, b sternal views).

Some commonly intercepted native and exotic species are illustrated and briefly described on the following pages.

Appendix A

Agrilus bilineatus (Weber):



Figure 4. *Agrilus bilineatus* (Weber)

- Pubescent line on pronotum extending down each elytron.
- Bronze-black to dark blue.
- Length 11 mm
- Host *Quercus* spp.

Agrilus cyanescens Ratzeburg



Figure 5 *Agrilus cyanescens* Ratzeburg

- No patches of pubescence.
- Metallic blue in color, darker below.
- Length 8 mm
- Host: *Lonicera* spp.
- **EXOTIC**

Agrilus anxius (Weber)



Figure 6. *Agrilus anxius* (Weber)

- No patches of pubescence.
- Bronze-black
- Length 12-13 mm
- Host: *Betula* spp. *Agrilus vittaticollis* (Randall)



Figure 7. *Agrilus vittaticollis* (Randall)

- Pubescent line on pronotum.
- Pronotum copper colored with dark elytra, darker below with violet highlights.
- Length 10-12mm
- Hosts: *Crataegus*, *Malus*, *Amelanchier* spp.

Appendix A

Agrilus obsoletoguttatus



Gory

Figure 8. *Agrilus obsoletoguttatus* Gory

- Pubescent spots on elytra.
- Copper color with violet elytral apices, darker below.
- Length 9-10 mm
- Hosts: Hardwood spp.

Agrilus masculinus



Horn

Figure 9. *Agrilus masculinus* Horn

- No distinct patches of pubescence
- Bronze-black, green face on males
- Length 6-7 mm
- Hindleg tarsi longer than tibia
- Host: *Acer negundo*

In addition to *Agrilus planipennis*, *A. subcinctus* Gory is the only other *Agrilus* species known to occur on *Fraxinus* in the Midwest. *A. subcinctus* differs behaviorally from *A. planipennis* in that it is a twig borer and prefers smaller branches for oviposition. In the field, *A. subcinctus* can be sweep-netted from ash leaves. *A. subcinctus* is

small in size (>4.0 mm) and features a distinct subbasal and subapical spots of scale-like pubescence on the elytra (Figure 10).



Figure 10. *Agrilus subcinctus* Gory

All of these *Agrilus* are often found on sticky and in funnel trap samples. With close to 50 *Agrilus* species known from the Midwest, I expect other species to be added to this preliminary guide in the near future.


Emerald Ash Borer Larval Screening Guide

J. E. Zablontny

Agrilus planipennis Fairmaire




A1-A6 Trapezoidal




A7 Bell-shaped

Agrilus anxius Gory



A1-A6 Rectangular



A7 Rectangular

The shape of the abdominal segments as well as host is useful for discriminating EAB (*Agrilus planipennis* Fairmaire) larvae from other native *Agrilus* species such as the bronze birch borer (*Agrilus anxius* Gory). I chose *A. anxius* to serve as a reference for comparison with EAB as its larval morphology is more typical for the genus.

The first abdominal segment shape varies considerably but can be distinctly rectangular to trapezoidal in *A. planipennis* while more rectangular and cylindrical in *A. anxius*. In EAB, A2 through A6 are somewhat trapezoidal with protruding flattened lobes. A7 is strongly bell shaped and A8 is somewhat trapezoidal in shape. In *A. anxius*, the abdominal segments do not have protruding flattened lobes and are rectangular in profile. A7 is not bell-shaped in bronze birch borer larvae.

These apparent differences in segment shapes are obvious in all instars of EAB examined so far. However, segment shape differences between species can be more subtle in prepupae and in poorly preserved larvae. In EAB prepupae, abdominal segment seven is still bell-shaped and rectangular in *A. anxius*.

In Michigan, EAB is the most common *Agrilus* species associated with *Fraxinus*. North American populations of EAB are known only to oviposit in *Fraxinus*.

Appendix C

CONTACTS

USDA APHIS PPQ

EAB Hotline: 1-866-322-4512 (Toll Free)

Paul Chaloux, EAB National Program Coordinator

Phone: 301-734-0917 e-mail: Paul.Chaloux@aphis.usda.gov

Philip Bell, Eastern Region EAB Program Manager, Raleigh, NC

Phone: 919-855-7312 e-mail: Philip.D.Bell@aphis.usda.gov

Roeland Elliston, Western Region EAB Program Manager, Fort Collins, CO

Phone: 970-494-7572 e-mail: Roeland.J.Elliston@aphis.usda.gov

Doug Bopp, Geographer, Brighton MI

Phone: 810-844-2727 e-mail: Douglas.A.Bopp@aphis.usda.gov

Ken Witt, Environmental Monitor, Brighton, MI

Phone: 810-8442716 e-mail: Kenneth.A.Witt@aphis.usda.gov

Sharon Lucik, Legislative and Public Affairs Specialist, Brighton, MI

Phone: 810-844-2713 e-mail Sharon.E.Lucik@aphis.usda.gov

James Buck, Ph.D., EAB National Program Analyst, Brighton, MI

Phone: 810-844-2733 e-mail: James.H.Buck@aphis.usda.gov


Identification Authority

James Zabloutny, Ph.D., USDA APHIS PPQ, Area Identifier

11200 Metro Airport Dr., ste. 140

Romulus, MI 48174

Phone: 734-942-9005 e-mail: James.E.Zabloutny@aphis.usda.gov



Cooperative Emerald Ash Borer Project



Information Packet

	<u>Toll Free Hotline</u>	<u>Website</u>
Illinois	800-641-3934	www.illinoiseab.com/
Indiana	866-663-9684	www.entm.purdue.edu/eab
Maryland	800-342-2507	www.mda.state.md.us/plants-pests/eab/
Michigan	866-325-0023	www.emeraldashborer.info
Ohio	888-644-6322	www.ohioagriculture.gov/eab
Pennsylvania	866-253-7189	http://www.dcnr.state.pa.us/forestry/fpm_invasives_EAB.aspx
West Virginia	866-322-4512	http://www.wvforestry.com/eab.cfm
Wisconsin	800-462-2803	dnr.wi.gov/org/land/Forestry/FH/Ash/
USDA	866-322-4512	www.aphis.usda.gov/ppq

For additional information you may also send email to the Cooperative Emerald Ash Borer Project at: EmeraldAshBorerInfo@aphis.usda.gov



Distinguishing Ash from other Common Trees

Diane Brown-Rytlewski and Rebecca Thompson
Michigan State University IPM Program

Identifying ash trees

Due to the recent discovery of the emerald ash borer (EAB) in Michigan, it is important to be able to recognize and identify ash trees. To date, emerald ash borer has only

been found on ash trees (see page 4). Not sure if your tree is an ash? This simple key is intended to help you distinguish between some common deciduous landscape trees frequently confused with ash, including: elm, boxelder, mountainash, walnut and hickory.

Using the identification key

Begin at number 1 on the key and choose (a) or (b). Then proceed to the number listed in italics at the end of your choice. This number will give you a new set of choices. Continue this way through the key. We have listed enough characteristics to help you determine whether or not your tree is an ash. If it doesn't match the characteristics in the key, relax; it most likely isn't an ash. If you don't want to use the

key, tree ID photos are on pages 2-4.

If you are still curious about what kind of trees you have, consult Extension Bulletin E-2332, "Identifying Trees of Michigan," or a good field guide such as: "The Tree Identification Book," by George W.D. Symonds; "Tree Finder: A Manual for the Identification of Trees by Their Leaves," by May T. Watts; or "A Field Guide to Trees and Shrubs" by George A. Petrides. There are many other good guides available; these are mentioned only as examples.

Identification key

1. a) Branches alternate (Fig. 1) – *go to 2*
b) Branches opposite (Fig. 2) – *go to 4*
2. a) Simple leaves, with irregular leaf base and toothed edge (Fig. 3). See **elm**, page 2.
b) Compound leaves (Fig. 4), with 9 to 15 leaflets, finely toothed around edge of leaf – *go to 3*
c) Compound leaves, 5 to 7 leaflets – *go to 3c*
3. a) Cut open twig lengthwise.
Chambered pith (Fig. 6). See **black walnut**, page 3.
b) If pith is not chambered, but has white flowers in May, orange or red berries in fall. See **mountainash**, page 3.
c) If pith is not chambered, but has three leaflets at end of leaf larger than the rest. See **hickory**, page 2.
4. a) Compound leaves, 5 to 9 leaflets, smooth or finely toothed around outer edge. See **ash**, page 4.
b) Compound leaves, 3 to 5 leaflets, few coarse teeth or none, end leaflet pointed (Fig. 5). See **boxelder**, page 4.



Fig. 1- Alternate branching



Fig. 2- Opposite branching



Fig. 3- Simple leaf (American elm)



Fig. 4- Compound leaf, 7 leaflets (White ash)



Fig. 5- Compound leaf, 3 leaflets (Boxelder)



Fig. 6- Chambered pith (Black walnut)

Line drawings: Steven Brown

Definitions

- Alternate – leaves/branches that are staggered or not directly across from each other, Fig 1.
- Opposite – leaves/branches that are directly across from each other, Fig. 2.
- Simple – a single leaf blade joined by its stalk to a woody stem, Fig 3.
- Compound – a leaf with more than one leaflet. All leaflets attached to a single leafstem, Fig. 4-5.

Appendix D


Elm, Ulmus spp.

Branches and buds are alternate and leaf bases are unequal. The leaves are simple, serrate (toothed) and 3 to 6 inches long and 1 to 3 inches wide (American elm) or 3/4 to 3 inches long and 1/3 to 1 inch wide (Siberian elm). The fruit is a winged seed.

Sandy Perry



Sandy Perry



Elm leaves and fruit



Unequal leaf base
Mary Wilson



Alternate branching


Left: American elm has a vase-shaped growth habit. Right: Siberian elms are fast-growing, brittle trees that break easily. Other elms will have similar leaves and seeds, although leaf size and growth habit will vary.

Hickory, Carya spp.


Shagbark hickory has distinctive bark – long, loose, shaggy strips. Leaves are compound, 8 to 14 inches long with 5 to 7 leaflets. The three terminal leaflets are larger than the other leaflets. Other similar species (not shown) include bitternut, pignut and mockernut, which have similar leaves and fruits although leaf size will vary. They do not have distinctive shaggy bark. These trees are seldom planted in landscapes, but are native, and may be found growing in woods. Fruits are hard-shelled light brown nuts, in a green husk that splits into four parts.

Hickory leaf

UGA0008353 Paul Wray



Bark of shagbark hickory tree
Diane Brown-Rytlewski



Hickory fruit

UGA0008213 Paul Wray



Appendix D



Black walnut fruit

UGA0008279 Paul Wray




Black walnut leaf


UGA0008448 Paul Wray

Black walnut, Juglans nigra

Branches and buds are alternate. Leaves are compound, 15 to 24 inches long, with 9 to 15 leaflets/leaf. Crushed leaflets and stems have a distinct odor – similar to turpentine. Twigs, split lengthwise, have chambered pith. Fruit is a large dark brown nut inside a green husk.



Black walnut chambered pith



Alternate branching

Mountainash — alternate branching


Diane Brown-Rytlewski



European mountainash, Sorbus aucuparia

Sorbus aucuparia

Leaves are compound, 5 to 9 inches long with 9 to 15 leaflets per leaf. Leaflets are a toothed, rounded oval shape. Flowers are five-petaled, white and similar to cherry or apple blossoms but in clusters. Fruits are fleshy, red-orange berries in clusters – they are found in the fall.



Mountainash leaf

Diane Brown-Rytlewski

Mountainash flowers

Diane Brown-Rytlewski

Appendix D

Ash, *Fraxinus* spp.

Branches and buds are opposite with a single bud at the end of the branch (terminal bud). Twigs are gray to brown and do not have a waxy coating. Leaves are compound, 8 to 12 inches long, 5 to 9 leaflets/leaf. Leaves may be finely toothed or have smooth edges. The most common ash trees planted in the landscape are **white ash** (*Fraxinus americana*) and **green ash** (*Fraxinus pennsylvanica*). Other native ash trees less commonly found include **black ash** (*Fraxinus nigra*) and

blue ash (*Fraxinus quadrangulata*) (not shown). **Black ash** has 7 to 11 leaflets and is found in wet woods; **blue ash** has 7 to 11 leaflets and distinctive 4-angled corky wings on the stem. **White ash** buds are paired with a leaf scar beneath the bud that looks like the letter “C” turned on its side. **Green ash** buds are paired with a leaf scar beneath the bud that looks like the letter “D” turned on its side (like a smile). Individual fruits are shaped like single wings and occur in clusters; many ash cultivars are seedless.




UGA0008289 Paul Wray

Green ash leaf



Diane Brown-Rytlewski

White ash leaf




Opposite branching

Dave Cappaert



Left: green ash leaf scar. Right: white ash leaf scar.

UGA0008169 Paul Wray




Ash seeds

Boxelder, *Acer negundo*


Boxelder is sometimes called ash-leaved maple. The twigs and buds are opposite; with a single bud on the end of the twig (terminal bud). Twigs are green to purplish brown, and often have a waxy white coating that can be rubbed off the

stem. Leaf scars beneath the buds are narrow, and join in a point. Leaves are compound, 4 to 10 inches long, with 3 to 5 leaflets/leaf. Leaves may have a few coarse teeth, or none. The end leaflet is sharply pointed. Fruit is a paired winged seed, occurring in clusters.



Boxelder leaf,
3 leaflets

UGA0008303 Paul Wray



Boxelder branches

UGA1219006 Bill Cook



Boxelder fruit

UGA1219003 Bill Cook

Photo credits:

Bill Cook, Michigan State University, www.forestryimages.org/

Paul Wray, Iowa State University, www.forestryimages.org/

MSU is an Affirmative-Action Equal-Opportunity Institution. MSU Extension programs and materials are open to all without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status, or family status. Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 20, 1914, in cooperation with the US Dept. of Agriculture. Margaret A. Bethel, Extension director, Michigan State University, E. Lansing, MI 48824.

The information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement or bias against those not mentioned. Reprinting cannot be used to endorse or advertise a commercial product or company.

Appendix D

Regulatory Information/Survey Form

Inspector Name: _____ Date: _____

Regulatory Contact

Concern or Business Name: _____

Contact Name: _____

Street Address: _____ City: _____

State: _____ Zip Code _____

County: _____

GPS Lat _____ Long _____

Phone: _____

Concern Type (circle up to three)

Nursery / Garden Center / Firewood Dealer / Public Campground / Private
Campground / Milling & Manufacturing / Landscaper / Research / Municipality /
Tree Removal Service / Landscaper / Logging Company / Misc. / Pallets / Pallet
Stock / Mulch

Aware of EAB Quarantine Y or N If yes,
how? _____

Interstate Movement Y or N

What/Commodity?

Frequency of shipments?

Quantity per shipment?

Final Destination of shipments?

From area near EAB positives?

Shipping Documents Available?

***Signature below confirms representative(s) of the Cooperative Emerald
Ash Borer Project contacted the signatory individual and gave them an
information packet about Emerald Ash Borer.***

Printed Name: _____

Signature: _____

Appendix D

Regulatory Information/Survey Form

Date: _____

Survey of Ash

Ash Product	Present? Y or N	Origin	Quantity	Inspected? Y or N	How long has the material been on site?
Firewood					
Nursery Stock					
Estab. Ash					
Ash Chips/Mulch					
Saw Logs					
Other					

Campground Survey


	Ash firewood Present? Y or N	% OH plates	% MI plates	% IN plates	Local Firewood Available	Permanent Campsites Present
Private Campground						
Public Campground						

EAB symptoms or specimens present Y or N,
If yes record ash product and X all indicators that apply.

Ash Product: _____

Crown Dieback	Life Stage	D-shaped holes	Bark splits	Epicormic shoots	Serpentine Gallery	Die-Back	Woodpecker Damage

Remarks:



Cooperative Emerald Ash Borer Project

Approximate
Range of ash species
in North America

Map reproduced from
USDA Forest Service source map




Cooperative Emerald Ash Borer Project

EAB locations in Illinois, Indiana, Michigan, Maryland, Missouri, Ohio, Pennsylvania, Virginia, Wisconsin, West Virginia and Canada


Appendix D

December 8, 2008



Map Key

- EAB positive
- Site under evaluation (symptoms, found in firewood, or eradicated)
- Federal EAB quarantine boundaries
- State quarantine generally treated area
- State quarantine
- State quarantine-other (SM)
- National Forests
- Canals and EAB regulated areas



Sources of available data:
 - Maryland Department of Agriculture
 - Michigan Department of Agriculture, Current Outbreak Details 2006-10-04
 - Michigan Department of Agriculture Statistics
 - Michigan Tech University School of Forestry
 - Illinois Department of Agriculture
 - Indiana Department of Natural Resources
 - Ohio Department of Agriculture
 - Pennsylvania Department of Agriculture
 - Virginia Department of Forestry
 - West Virginia Department of Forestry
 - Emerald Ash Borer Survey Results as of October 10, 2008
 - USGS National Wetlands Inventory
 - USGS National Wetlands Inventory
 - USGS National Wetlands Inventory
 - USGS National Wetlands Inventory



US Department of Agriculture
 Animal and Plant Health Inspection Service
 Forest Protection Service

UCDAP #10-000
 Map by S. Kopp

Guidance on Conducting Select Regulatory Activities in the Emerald Ash Borer Program

INTRODUCTION

Initiation of the National Emerald Ash Borer (EAB) Survey in 2008 will involve numerous additional states in Program activities, potentially including regulatory activities. This document is intended to provide some guidance on conducting a sampling of regulatory activities in the EAB Program, with an emphasis on those aspects of regulatory work that are unique to the Program. It is not intended to be inclusive or to serve as a manual for setting up and running a regulatory section in support of the EAB Program.

REGULATORY AUTHORITIES

In October 2003 APHIS issued a Domestic Quarantine Notice, 7 CFR 301.53 (1)-(9), establishing the emerald ash borer Quarantine. This quarantine defines the regulated area and provides the authority to conduct regulatory activities. The focus of USDA APHIS PPQ regulatory activities fall under 7 CFR.301.53 (4)-(9) which set the conditions governing interstate movement of regulated articles. Several revisions to this quarantine have been published as interim rules or Deputy Administrator issued letters to the State Plant Regulatory Officers. Broad-reaching Federal authority for invasive pests and plant protection also resides in The Plant Protection Act of 2000.

REGULATED ARTICLES

7 CFR 301.53-2(a) classifies the following as **regulated articles**:

- The emerald ash borer in any stage of development
- Firewood of all non-coniferous (hardwood) species
- If in the genus *Fraxinus*
 - nursery stock,
 - green lumber,
 - and other material living, dead, cut, or fallen, including logs, stumps, roots, branches, composted and uncomposted chips of the genus *Fraxinus*.

Any other article, product, or means of conveyance not listed in 301.53-2(a) may also be designated as a regulated article if determined by the EAB Program that it presents a risk of spreading EAB in accordance with 301.53-2(b). In 301.53-1 an infestation is defined as the presence of EAB or the existence of circumstances that make it reasonable to believe that EAB is present. This definition, in conjunction with 301.53-2(b), will allow the EAB Program to take certain actions as required to regulate interstate movement of articles not otherwise covered, or in circumstances not covered by the quarantine. Intrastate movement of regulated articles is addressed under state authorities.

REGULATED AREAS

Federal Quarantines

A map of the areas under federal quarantine is available at the following website:

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/downloads/multi-stateeab.pdf

State Quarantines

Several states have also established intrastate quarantines for EAB. These are usually on a county-by-county basis. Please consult the office of the State Plant Regulatory Official for the state in questions for specifics on state quarantines and regulations. Contact information for State Plant Regulatory Officials can be found at the following web site:

<http://www.nationalplantboard.org/member/index.html>

COMPLIANCE AGREEMENTS

Entering into compliance agreements with establishments engaged in regulated activities allows the program to make efficient use of regulatory personnel. The EAB Program currently enters into compliance agreements with establishments engaged in the following activities:

- Firewood producers
- Firewood distributors
- Sawmills that handle non-coniferous wood
- Transporting of non-coniferous timber and/or green lumber
- Production of mulch and/or ash chips
- Treatment of wood products or packing materials that contain ash

Samples of compliance agreements used by the program can be found at the following website:

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/regulatory.shtml

Prior to entering into a compliance agreement with an establishment or business, a site visit to determine ability to meet the terms of the compliance agreement must be made by an authorized inspector. Those requirements must be explained to the establishment, along with the consequences for non-compliance. Signing of the compliance agreement should occur only after all parties are satisfied that the terms of the compliance agreement can and will be met.

Compliance Agreement Monitoring

After a compliance agreement is entered into, periodic visits to monitor continued adherence to requirements must occur. The frequency of these visits is dependent upon a number of factors. The risk assessment matrix below is an example of a tool developed to quantify the risk level created by a regulated establishment, and the consequent minimum period that should elapse between compliance monitoring visits.

Risk Assessment Matrix for EAB

Below are descriptions of risk factors used in the risk assessment matrix to calculate the frequency of compliance monitoring visits to establishments operating under federal compliance agreements. An inspector’s judgment is an important tool in determining risk, and other factors not listed below also may be considered.

Risk Factors:

- Ease of Establishment of EAB - Determine potential risk of establishment of EAB given state of pest and condition of host (nursery stock would be highest risk)
- Traceability – (inverse risk) if the business operation has easily followed paper trails then the risk is low
- Profile of the Concern – (inverse risk) high profile = low risk. Degree to which concern moves regulated material in large marked vehicles and operates in the public arena.
- Location of Activity –use following zones for risk assignment:
 - High = Within 50 miles of the quarantine boundary
 - Medium = buffer zone within quarantine boundary (50 to 100 miles from boundary)
 - Low = within quarantine area more than 100 miles from boundary
- Mobility of the Regulated Article – The more mobile the higher the risk

Calculating Risk Level: *assign level to each risk factor and multiply times the given factor for total*

Add Total column to arrive at quantified risk level

Example 1: Big nursery operation 55 miles from quarantine boundary

Risk Level = 9

RISK LEVEL	Ease of Establishment of EAB	Traceability	Profile of Concern	Location of Activity	Mobility of Regulated Articles	Total
LOW (x1)		X	X			2
MEDIUM (x2)				X	X	4
HIGH (x3)	X					3

Example 2: Low profile firewood dealer in northern gateway
 Risk Level = **14**

RISK LEVEL	Ease of Establishment of EAB	Traceability	Profile of Concern	Location of Activity	Mobility of Regulated Articles	Total
LOW (x1)						0
MEDIUM (x2)	X					2
HIGH (x3)		X	X	X	X	12

Example 3: USFS Campground 60 miles from quarantine boundary
 Risk level = **11**

RISK LEVEL	Ease of Establishment	Availability of Records	Profile of Concern	Location of Activity	Mobility of Regulated Articles	Total
LOW (x1)						0
MEDIUM (x2)	X	X	X	X		8
HIGH (x3)					X	3

Use the following levels to assign a priority for the regulatory check data base
 Risk Level 5-6 = Priority 4 – visit as needed
 Risk Level 7-9 = Priority 3 – visit semi-annually
 Risk Level 10-12 = Priority 2 – visit bi-monthly on a minimum
 Risk Level 13 – 15 = Priority 1 – visit monthly on a minimum

Firewood Quality Assurance Plan

The firewood quality assurance plan below provides specific guidance on the types and frequency of compliance monitoring activities used for firewood production and distribution.

Purpose: To increase firewood producer accountability and monitor alignment with compliance agreement parameters beyond initial point of production.

Firewood Compliance Agreement Monitoring
 Base Line Periodicity

Firewood Treatment	Monitoring Guidance
Debarking and 1/2" of wood removed	Weekly during firewood season

Heat Treatment, Fumigation, Kiln Sterilization	Weekly during firewood season
Safeguarding (i.e. Distribution Facilities)	Monthly

Monitoring Production Facilities

1. Monitoring includes site visits, product inspection, and equipment data review.
2. Random visits should be conducted during non-production season to verify status.
3. The following information should be reviewed/collected periodically as a paper audit and to provide a basis for monitoring distribution facilities.
 - a. Customer List
 - b. Invoiced quantities
 - c. Shipment dates
 - d. Production dates
4. Discussions should be held with staff to determine their degree of knowledge regarding EAB regulated items and their handling.

Monitoring Distribution Facilities

1. Monitoring should occur at first transportation event and a minimum of monthly thereafter.
2. A minimum of 10% of the product should be inspected to assure labeling and/or mitigation/treatment requirements are met.
3. The following information should be reviewed/collected periodically as a paper audit.
 - a. Customer List
 - b. Invoiced quantities
 - c. Shipment dates
 - d. Production/supplier information
4. If a shipment has an interstate destination, it is recommended to randomly notify the destination state SPHD and request them to do a quality assurance follow-up check at product destination.
5. Discussions should be held with staff to determine their degree of knowledge regarding EAB regulated items and their handling.

Monitoring Firewood in the Marketplace

1. It is recommended that PPQ regulatory officers in each state make it a routine practice to monitor firewood in retail establishments to determine if the product is regulated.
2. If the firewood is regulated it should be inspected to determine compliance with federal regulations regarding that product’s movement.
3. It is suggested that at least 5 of these checks are conducted per month.
4. These quality assurance monitoring events should be used as an opportunity to provide outreach and education materials to the retail establishment for potential posting in facility retail or employee areas.

Mulch Sampling Protocol

Size Requirements and Screening Procedure for Hardwood Mulch and Chips in the Emerald Ash Borer Quarantine Area

The following procedure should be followed to determine if hardwood and bark chips, nuggets, and mulch materials can be considered safe for movement from the emerald ash borer quarantine area:

- Step 1. Using a 12 inch diameter 3.25 inch deep sieve with 1.25 inch steel mesh openings (Fisher Scientific #04-884-1J) take 10 samples from random locations in the chip or mulch pile - - do not take all samples from the same location. If any chips are found that are greater than 2.5 inches in two dimensions the pile is rejected. If there are no chips found greater than 2.5 inches in two dimensions then proceed to Step 2.
- Step 2. If four or more chips from the 10 samples do not pass through the sieve proceed to Step 3. If three or fewer chips from the 10 samples do not pass through the sieve, then the pile passes and can be moved.
- Step 3. Resample. Take 10 additional samples from random locations in the chip pile. If any chips are found that are greater than 2.5 inches in two dimensions the pile is rejected. If there are no chips found greater than 2.5 inches then proceed to Step 4.
- Step 4. If four or more chips from the 10 samples do not pass through/put through the sieve the pile is rejected. If three or fewer chips fail to pass through the sieve then the pile passes and can be moved.

Note: Mulch chips that do not meet the specified requirements as outlined above may be reground to meet this specification of 1 inch in two dimensions or alternatively composted as outlined below.

Composting Option For Mulch That Fails Sampling Protocol

Composting Requirements for Hardwood Mulch and Chips in the Emerald Ash Borer Quarantine Area

To render hardwood and bark chips, nuggets, and mulch materials that are larger than 1.0 inches in two dimensions certifiable/safe for movement they can be composted using the following procedure:

- Step 1. Compost piles must be a minimum of 200 cubic yards.
- Step 2. Internal temperature at a depth of 18 inches must reach 140 °F (60 °C) for 4 continuous days.
- Step 3. Using a front-end loader or a bulldozer, remove the outer layer of the compost pile to a depth of 3 feet.
- Step 4. Start a second compost pile using the recently-removed cover material as a core.
- Step 5. Move the core material from the first compost pile and place on the second compost pile as a cover at least 3 feet deep.

Step 6. Allow the second compost pile to remain undisturbed until the temperature reaches 140 °F (60 °C) for at least 4 continuous days.

Step 7. Remove the second compost pile and use as fully-composted material.

This procedure will allow continuous operation. After the first compost pile is “turned” to become the second compost pile, a new “first” compost pile can be started.

LIMITED PERMITS AND FEDERAL CERTIFICATES

Limited Permits and Federal Certificates are instruments used to allow the movement of regulated articles out of quarantined areas. Limited Permits allow for movement to a restricted set of locations, often a single location. Restrictions on movement are detailed on the limited permit issued to the article in question. Federal Certificates allow regulated articles to move to any destination in the United States. Both instruments will contain additional details on the mode of transport and the type and quantity of the article being moved.

Common forms of the Limited Permit include PPQ Form 530, PPQ Form 537 (self-adhesive labels), pre-printed labeling on packaged goods and electronic stamps. In all cases, the Limited Permit remains the property of USDA APHIS. In the EAB Program, PPQ Forms 530 and 537 are used and application of the Limited Permit is restricted to ash logs, lumber, stumps, branches and chips being transported to an approved treatment facility located outside the quarantined area. Both the transporter and the treatment facility must operate under a compliance agreement with the program. Specific conditions governing this movement are contained in the compliance agreement “Transport of Regulated Articles”. Limited permits are also issued to non-compliant commodities being returned to the state of origin during the course of regulatory actions taken in response to quarantine violations.

Federal Certificates can be issued in several forms as well, including PPQ Form 540, pre-printed packaging materials and electronic stamps. The EAB Program uses PPQ Form 540 as well as rubber and electronic stamps. Regulated articles eligible for interstate movement when accompanied by a Federal Certificate include ash lumber and logs, stumps and branches, or chips and firewood after they have been processed or treated at a facility operating under a compliance agreement. Specific conditions governing movement of these articles are contained in the compliance agreement relating to the article in question.

Prior to issuance of a Limited permit or Federal Certificate, eligibility of the article for interstate movement must be assured. This assurance is accomplished by entering into a compliance agreement with the person or establishment moving the article, or by the on-site presence of an authorized inspector. If the latter situation occurs, the inspector is responsible for ensuring that all necessary conditions have been met, and for signing and issuing the Limited Permit or Federal Certificate.

SPECIAL OPERATIONS

The EAB Program conducts special operations (sometimes referred to as blitzes) to serve as deterrents and quality control checks for movement of regulated articles. These operations are conducted in concert with state cooperators or less frequently as stand alone federal operations. Violations are issued and investigatory procedures are invoked when it is appropriate to do so for any individuals found moving regulated articles in violation of the federal quarantines. Violations of intra-state quarantines are referred to appropriate state regulatory officials.

Procedural details for planning and conducting a roadside firewood operation are given below as an example:

Roadside Firewood Operations

Early Planning Guidelines

1. **Cooperators:** Define the cooperators to participate in the operation and meet with them well in advance of the operations. Clearly define each others roles and authorities especially as it applies to stops, searches, seizures and what to do with individuals who do not cooperate. Set up a contact point person for each cooperator in the operation. Cooperators to consider are:
 - a. State Dept. of Agriculture- This is the primary cooperator and operations should be staffed with both federal and state regulators. Identify those people who can document the federal safeguarding action/violation either on EAN (PPQ523) or ROV (PPQ 518) forms (validity of cooperator cards as authority to issue violations appears to be in question by OGC). Identify who can issue paperwork for state violations. Working in (state/federal) pairs at the pull-off can help the operation run more smoothly and get violators back on the road quickly.
 - b. Law Enforcement – Law enforcement cooperators typically include state police for major routes, sheriffs for secondary roads, and municipal police within city/town limits. Motor carrier enforcement is not typically involved in firewood operations; they are more often cooperators when monitoring log movement. Law enforcement generally provides a squad car parked with flashing lights just prior to the stop and a uniformed officer for support. We have found that this is a beneficial arrangement as law enforcement often will find violations of their own or the officers themselves become engaged and participate in the outreach. Also pay attention to jurisdiction—whether the law enforcement branch you go with has the right to exercise the law at your particular site. For example, a county sheriff’s deputy in one state related that State Highway Patrol can catch violations on state highways, but once the person is on private property (i.e. the pull off) their jurisdiction ends. A potential benefit for State Departments of Agriculture you may want to explore is determining if the salary/costs of the officer qualify as part of cost sharing towards a federal dollars match in an EAB cooperative agreement. *Note: County sheriffs (and a squad car) are generally less expensive and more available during high-traffic weekends like Memorial/Labor Day. If the*

- squad car with flashing lights is parked on the side of the road, have the uniformed officer up at the pull-off to provide security for workers. Violators are much less likely to react aggressively with an (armed) officer close by.*
- c. State DOT- DOT should be contacted to discuss the operation site and provide traffic control equipment to maximize public and officer safety. Equipment usually includes a flashing marquee about ½ mile before the stop, traffic cones, and barrels. It's a good idea to put "Road Work Ahead" signs for the oncoming lanes so that drivers headed opposite the road block aren't surprised and have a chance to slow down.
 - d. Public Information Staff – Publicizing these events is critical to continued public education and outreach, but when and how is state driven. In some states they must publicize a roadside stop prior to conducting it. In those states a generic notification has been sent out saying firewood operations would occur over the weekend, but not detailing exactly when or where. Several operations have also invited media to the event itself where they are allowed to shoot footage and usually the PIO meets them there to answer questions and provide outreach. One operation was even published in USA Today.
 - e. APHIS-IES- It is beneficial to involve IES in the early planning. They may opt not to be present for the operation but as they will be receiving the Notifications of Violation, they should participate in set-up.
 - f. Native Americans - Any state conducting operations with federally recognized Native American groups may want to outreach to them to let them know and have a voice as appropriate.
 - g. Misc. – Other state agencies may want to participate such as the DNR. Incorporate these groups as you see fit. If the operation is done close to a state line, the adjoining state should be notified.
2. **Site Selection:** Careful attention should be made to site selection addressing all of the following factors at a minimum. Always secure permission from the landowner (public or private) before choosing a site and be sure to send them a thank-you note after the operation has concluded especially if you'd like to use the site again in the future. If you are not familiar with an area you'd like to blitz, call local law enforcement and DOT garages for suggestions. Sometimes the "perfect" site is not safe for an operation and these professionals can help make that call.
- a. Potential Firewood Movement – Evaluate the route on which the operation is proposed. Does the route provide relatively direct access to recreational areas where camping will occur or is it an artery into the state for vacationers? Will travelers likely be coming from an infested/regulated area? Is it near posted signs that warn motorists they are crossing a quarantine line? Are there firewood dealers nearby inside the quarantine that are selling wood (legally) without informing customers of quarantine laws?
 - b. Traffic Volume – At this point, no mandatory stop operations have been conducted on interstate highways for valid reasons. Volume of traffic is high, vehicle speeds are high, and stop logistics untenable for law enforcement. Roadside rest areas can be used for "voluntary" firewood checkpoints. Provide an electronic sign so that drivers exit the freeway and conduct the operation in the

- parking lot. The best options thus far have come from secondary US routes and state routes that are two lane roads. It is extremely useful to place the stop at an area where the road widens into 3 lanes for a short time. Officers can generally work safely in the middle lane. On these secondary routes traffic volume is sufficient to make operations effective but, not such a great volume that officers are creating back-ups. One consideration we have agreed to with law enforcement is to stop operations if the back-up becomes too great and allow a period of free movement. You may also opt to wave commercial/municipal vehicles through the block to keep traffic flowing at a reasonable rate.
- c. Physical Conditions – The site must provide regulators the ability to stop cars safely, provide an inspection area, route vehicles back into traffic, and provide staff parking. Examples of sites used have included: two lane road near church with church’s permission to use parking lot for inspections and church bathroom for officers; three lane route with temporary storage yard permission to use parking lot for inspections, gas stations nearby; two lane route with extensive shoulder/parking area in front of residence/business with owner permission; two lane route with historic marker pullout for inspection; town street with Elks Lodge permission to use parking lot for inspection and lodge for bathrooms (although they were not too happy with us when they realized the town cop would be with us all day watching who came in to drink and how long they were there).
 - d. Amenities – Operations vary in length from 6 to 12 hours typically and needs of the regulatory staff must be considered. Restroom facilities must be provided within a reasonable distance. If no eating facilities are near-by staff needs to be informed so they can prepare for that. The program has several shade tents and usually sets one up to give officers a break from the sun and/or rain. It can be very hot standing on the tarmac in the sun. The program has provided ice chests and management has provided bottled water and ice (from our own pockets-not govt. approved). Chairs and a folding table are also a real benefit to allow people breaks. This is no place to work on a suntan—sun block and appropriate clothing is essential, even on cloudy days.
 - e. Firewood Disposal – This is a critical component. If the operations are conducted outside a quarantined area (usual circumstance) then the material collected must be safeguarded or treated immediately. To date the program has utilized enclosed vehicles to store wood and transport it back into the quarantined area or burned it expeditiously on site or near-by. Another option would be to have a chipper on site and chip the material to less than 1” in 2 dimensions. Some state run operations have routinely allowed the individuals to burn the material within 48 hours. The EAB program does not support that disposition option. *Note: When disposal issues can not be resolved, an alternative approach is to set up a road block just inside the quarantine boundary and warn motorists that they are about to break the law. This is particularly useful in areas where the quarantine has just been expanded/imposed and serves as more of an outreach than regulatory activity*
3. **Operation Date:** The most effective dates for operations are those that precede some event in the area that draws the camping public. This can be as obvious as traditional

holiday weekends including Memorial Day, Labor Day, and 4th of July. There are many other events that may occur in your area that draw large crowds that camp. Check state park event listings for other ideas. Remember, camping season isn't just between Memorial and Labor Days—people tend to bring more firewood for the chilly fall nights. Some we have run across are:

- Bluegrass music festivals
- NASCAR races
- Coon dog trials
- Native American pow-wows
- Fall color tourism
- Hunting season
- Halloween (a large camping draw in Ohio)
- Horse camping events
- Fishing tournaments
- Morel season
- College rivalry sporting events

4. **Equipment:** What follows is a list of equipment the program has found useful at operations.
 - a. Shade tent
 - b. Folding table
 - c. Folding chairs
 - d. Orange pylons
 - e. Cooler
 - f. Enclosed vehicle for firewood transportation
 - g. Operations Box (OpsBox) – plastic tubs w/lid kept in readiness
 1. Work gloves
 2. Vials with alcohol & larval forceps
 3. Chisels
 4. Outreach material – pamphlets, tattoos, EAB ID guides, etc. Bring hundreds of brochures—this may be your only chance to get EAB materials directly into the hands of these people.
 5. Accordion folders with: operation protocol, current quarantine maps, current quarantine language, EANs, PPQ 518s, maps to closest hospital/urgent care facility, PPQ 391s, directions to disposal site if applicable, lists of participants and contact numbers, operation log, survey forms, and written documentation of your agency's authority to conduct this operation and seize private property
 6. Hand held STOP/SLOW signs
 7. Safety vests – reflective mesh with pocket
 8. Orange flags
 9. First aid kit
 10. Flashlight
 11. Digital camera
 12. Outreach folder—violators are often angry and complain that they've never even heard of EAB or firewood movement. Compile a binder/booklet/file of local newspaper articles, TV station transcripts, pictures, posters, etc. to avoid

unpleasant phone calls to government offices made by those inconvenienced people.

Firewood Operation Staffing Duties

Roadway

Law Enforcement: Slow traffic flow ahead of the blitz

Vehicle Profiler (2):

- Tallies vehicle numbers
- Assesses vehicle for secondary referral
- Marks referred vehicle with flagging tape or radios ahead
- Provides EAB literature to drivers waved through

Flagger (1):

- Directs cars into inspection area and out of inspection area

Inspection Area

Interviewer/Inspector (4 – two teams):

- Greet and present identification
- Ask drivers if they are carrying firewood
- Ask about carrying firewood and ask to inspect the vehicle
- If no firewood found thank the driver, provide EAB pamphlet, remove flagging, and direct back into traffic
- If illegal hardwood firewood present seize and safeguard, explain why
- firewood is illegal, complete seizure record and a PPQ 523 EAN,
- take pictures of firewood with an identifier in the picture
- Inform the driver about penalty procedures

Other

Staff to alternate with other positions (1-2)

Supervisor (minimum 1)

Conducting Roadside Firewood Operations

- Personnel should be properly dressed according to their agency guidelines but at least wearing or carrying an item that identifies them as regulatory officials. This is for safety and program identification.
- It is recommended that each inspector have a flashlight for inspection after dark and a pair of gloves for unloading seized firewood.
- Law enforcement involvement is mandatory. In most scenarios law enforcement should be present and participating. If on site law enforcement is not possible it is still mandatory to contact local law enforcement and inform them about the operation

and solicit their support. A list of appropriate law enforcement contact numbers must be on site as well as a map to the nearest trauma facility. Traffic management should be handled by law enforcement and the use of safety aids should be maximized; electronic signs, reflective cones, reflective vests, etc.

- Traffic will be slowed, preferable by law enforcement, and regulatory personnel will visually inspect vehicles for evidence of regulated materials. If no regulated materials are present, thank the motorist for their cooperation, provide an educational pamphlet, and waive these vehicles on. When regulated materials are present, or there is an indication that regulated materials may be present as evidenced by related materials, the vehicles will be directed to the designated secondary inspection areas. All secondary inspections need to take place out of the flow of traffic in an area safe for travelers to enter/exit vehicles and for thorough inspection to take place. Cones should be used to mark of inspection areas and promote safe traffic flow.
- Contact with vehicles and travelers referred for secondary inspection should be conducted safely and with the following points in mind:
 - Do not approach a vehicle alone, work in pairs
 - Do not enter a vehicle or trailer unless another inspector is aware of your intent and is monitoring the situation
 - If approaching a vehicle at night, use a flashlight
 - Visually check the car and occupants for weapon or hazardous situations before focusing on the survey interview
 - Be aware of vehicle occupants as you talk with them
 - Never stand directly in front of a vehicle
 - If violence is encountered, retreat to a safe place if possible, contact and wait for law enforcement
 - In a polite and professional manner, identify yourself, present badge/identification and explain the reason for your contact, explain the program and its purpose, and the programs impact and threat to the nation's resources. Describe the Quarantines (have copies available) and explain the ban on movement of regulated materials from the regulated areas.
 - When a motorist is found transporting regulated materials, conduct an interview gathering information expeditiously.
 - If regulated material is found moved in violation of the quarantine, regulatory action will be taken. Seizure of all of non-compliant regulated materials is required. Explain the enforcement activity and quarantine. e Plant Protection Act of 2000 and appropriate state legislations
 - A PPQ 523 (Emergency Action Notification) shall be completed for each regulatory seizure.
 - For all commercial violations contact a supervisor.
 - Confiscated regulated materials should be seized and safeguarded in a safe and efficient manner.
 - Should a motorist become belligerent and refuse to give up their fire wood or attempt to "run" the checkpoint remove yourself to a safe place and contact law enforcement. Do not attempt to detain travelers for non-compliance. Despite their non-compliance,

gather as much information as possible and enter it into the survey data, especially the license plate number. Further regulatory follow up will be required.

- Enclosed vehicles or containers must be on hand to safeguard confiscated items for the duration of the operation. The operation site may not be vacated until all confiscated materials are safeguarded or removed for destruction.

Remember, you are representing the United States Departments of Agriculture. Many citizens will be transiting through the operation area. Present a professional and informed presence. Use good common sense and, most importantly: **WORK SAFE!!!!**

Other special operations conducted by the EAB Program include weigh station and highway rest area operations and campground operations. Weigh station operations typically target logging trucks moving logs out of quarantine. Rest area operations focus on motorists moving firewood and also provide excellent outreach opportunities. Campground operations target campers moving firewood and also provide outreach opportunities.

VIOLATIONS/INVESTIGATIVE AND ENFORCEMENT SERVICES

When individuals or companies violate federal regulations their actions may result in the case being turned over to Investigative and Enforcement Services (IES). EAB Program officers will issue Limited Permits to non-compliant commodities when necessary and fill out required paperwork, Emergency Action Notification and Notices of Violation, as described in the Manual for Agricultural Clearance. Officers will also provide statements that are complete and accurate and suitable for an affidavit. EAB officers are responsible to identify and secure for the investigator all the documents, items or photographs that may be used as evidence. Chains of custody will be initiated by program officers for items of evidence such as EAB specimens, ash firewood or shipment paperwork if necessary.

Non-Commercial Violations

The movement of firewood presents a significant risk for artificial spread of EAB. This movement has a commercial component but a large percentage of it is done by individuals who are moving it for personal use to go camping, supply vacation homes, or provide fuel for winter heating. The EAB Program has made a concerted effort to educate this segment of the population about EAB and the risks in moving firewood. This is particularly difficult in areas where woodlots abound and camping and hunting are major recreation activities. The program soon realized that an EAB educated but non-compliant portion of the public continues to move firewood. It has been the past practice of IES and PPQ to not penalize non-commercial violations of domestic quarantines. This past practice gave the program no appropriate deterrent for these violations. EAB program officials worked with IES and CPAT to create a stronger regulatory platform for the program. It was agreed that the non-commercial violations would be reviewed by IES field investigators, but not investigated, and then referred to headquarters. IES staff in headquarters assesses a penalty using a newly revised table of penalties (attached in

appendix V). In general, stipulations begin at \$250.00 per non-commercial violation but may be higher depending on aggravating factors.

Commercial Violations

Commercial violations of the EAB regulations are processed through IES as in the past. Notices of Violation, associated paperwork and evidence is forwarded to IES field staff for further investigation, then submission to headquarters.

Cooperation with State Regulatory Enforcement

When less than an entire state is quarantined program officers may encounter violations of the state quarantine that do not involve interstate movement. 301.53-1 defines an infestation to be the presence of EAB or the existence or circumstances that make it reasonable to believe that EAB is present. This stipulation, along with 7 CFR 301.53-2(b) stating that any other article, product, or means of conveyance not listed in 301.53-2(a) may also be designated as a regulated article if determined by the EAB Program that it presents a risk of spreading EAB, will allow officers to take appropriate safeguarding action in this situation. The program officer should immediately make contact with a state regulatory official and fill out a PPQ 523 (EAN) if necessary. The officer should prepare a statement for use by state officials.

Guidelines for Use in Preparing to do a Firewood Operation

The following items may be useful to consider when planning to do a firewood operation roadside stop.

1. Cooperators

Define the cooperators to participate in operation and meet with them well in advance of the operations. Clearly define each others roles and authorities especially as it applies to stops, searches, and what to do with individuals who do not cooperate. Set up a contact point person for each cooperator in the operation. Cooperators to consider are:

- a. State Dept. of Agriculture - This is the primary cooperator and operations should be staffed with both federal and state regulators.
- b. Law Enforcement – Law enforcement cooperators typically include state police for major routes, sheriffs for secondary roads, and municipal police within city/town limits. Motor carrier enforcement is not typically involved in firewood operations; they are typically cooperators monitoring log movement. Law enforcement generally provides a squad car parked with flashing lights just prior to the stop and a uniformed officer for support. We have found that this is a beneficial arrangement as law enforcement often will find violations of their own, or it is OT for the officer, or they officers themselves become engaged and participate in the outreach. A benefit for State Departments of Agriculture you may discuss using the salary/costs of the officer as part of cost sharing towards federal dollars match.
- c. State DOT- DOT should be contacted to discuss the operation site and provide traffic control equipment to maximize public and officer safety. Equipment usually includes a flashing marquee about ½ mile before the stop, cones and barrels.
- d. Public Information Staff – Publicizing these events is critical to continued public education and outreach, but when and how is state driven. In some states they must publicize a roadside stop prior to conducting it. In those states a generic notification has been sent out saying firewood operations would occur over the weekend, but not detailing exactly when or where. Several operations have also invited media to the event itself where they are allowed to shoot footage and usually the PIO meets them there to answer questions and provide outreach. One operation was even published in USA Today.
- e. APHIS-IES - It is beneficial to involve IES in the early planning. They may opt to be present for the operation or not, but as they will be receiving the Notifications of Violation, they should participate in set-up.

- f. Native Americans - Any state conducting operations with federally recognized Native American groups may want to outreach to them to let them know and have a voice as appropriate.
- g. Misc. – Other state agencies may want to participate such as the DNR. Incorporate these groups as you see fit. If the operations is done close to a state line, the adjoining state should be notified.

2. Site Selection

Careful attention should be made to site selection addressing all of the following factors at a minimum.

- a. Potential Firewood Movement – Evaluate the route on which the operation is proposed. Does the route provide fairly direct access to recreational areas where camping will occur or is it an artery into the state for vacationers? Will travelers likely be coming from an infested/regulated area?
- b. Traffic Volume – At this point, no mandatory stop operations have been conducted on interstates for valid reasons. Volume of traffic is high, vehicle speeds are high, and stop logistics untenable for law enforcement. The best options thus far have come from secondary US routes and state routes that are two lane roads. It is extremely useful to place the stop at an area where the road widens into 3 lanes for a short time. Officers can generally work safely in the middle lane. On these secondary routes traffic volume is sufficient to make operations effective but, not such a great volume that officers are creating back-ups. One consideration we have agreed to with law enforcement is to stop operations if the back-up becomes too great and allow a period of free movement.
- c. Physical Conditions – The site must provide regulators the safe ability to stop cars safely, provide an inspection area, route vehicles back into traffic, and provide staff parking. Examples of sites used have included: two lane road near church with church's permission to use parking lot for inspections and church bathroom for officers; three lane route with temporary storage yard permission to use parking lot for inspections, gas stations nearby; two lane route with extensive shoulder/parking area in front of residence/business with owner permission; two lane route with historic marker pullout for inspection; town street with Elks Lodge permission to use parking lot for inspection and lodge for bathrooms (although they were not to happy with us when they realized the town cop would be with us all day watching who came in to drink and how long they were there).
- d. Amenities – Operations vary in length from 6 to 12 hours typically and needs of the regulatory staff must be considered. Restroom facilities must be provided within a reasonable distance. If no eating facilities are near-by staff needs to be informed so they can

prepare for that. The program has several shade tents and usually sets one up to give officers a break from the sun and/or rain. It can be very hot standing on the tarmac in the sun. The program has provided ice chests and management has provided bottled water and ice (from our own pockets-not govt. approved). Chairs and a folding table are also a real benefit to allow people breaks.

- e. Firewood Disposal – This is a critical component. If the operations are conducted outside a quarantined area (usual circumstance) then the material collected must be safeguarded or treated immediately. To date the program has utilized enclosed vehicles to store wood and transport it back into the quarantined area or burned it expeditiously on site or near-by. Another option would be to have a chipper on site and chip the material to less than 1" in 2 dimensions. Some state run operations have routinely allowed the individuals to burn the material within 48 hours. The EAB program does not support that disposition option.

3. **Operation Date**

The most effective dates for operations are those that precede some event in the area that draws the camping public. This can be as obvious as traditional holiday weekends including Memorial Day, Labor Day, and 4th of July. There are many other events that may occur in your area that draw large crowds that camp. Some we have run across are:

- Bluegrass music festivals
- NASCAR races
- Coon dog trials
- Native American pow-wows
- Fall color tourism
- Hunting season
- Halloween (a large camping draw in Ohio)
- Horse camping events
- Fishing tournaments
- Morel season
- College rivalry sporting events

4. **Equipment**

What follows is a list of equipment the program has found useful at operations.

- a. Shade tent
- b. Folding table
- c. Folding chairs
- d. Orange pylons
- e. Cooler

- f. Enclosed vehicle for firewood transportation
- g. Operations Box (OpsBox) – plastic tubs w/lid we keep ready to go
 - a. Work gloves
 - b. Vials with alcohol & larval forceps
 - c. Chisels
 - d. Outreach material – pamphlets, tattoos, EAB ID guides, etc.
 - e. Accordion folders with: operation protocol, current quarantine maps, current quarantine language, EANs, PPQ 518s, maps to closest hospital/urgent care facility, PPQ 391s, directions to disposal site if applicable, lists of participants and contact numbers, operation log and survey forms
 - f. Hand held STOP/SLOW signs
 - g. Safety vests – reflective mesh with pocket
 - h. Orange flags
 - i. First aid kit
 - j. Flashlight
 - k. Digital camera

Firewood Operation Staffing Duties

Roadway

Law Enforcement: Slow traffic flow ahead of the operation

Vehicle Profiler (2):

- Tallies vehicle numbers onto log sheet
- Assesses vehicle for secondary referral
- Marks referred vehicle with flagging tape or radios ahead
- Provides EAB literature to drivers waved through

Flagger (1):

- Directs cars into inspection area and out of inspection area

Inspection Area

Interviewer/Inspector (4 – two teams):

- Greet and present identification
- Ask drivers where they are coming from and if they are carrying firewood
- Ask to inspect the vehicle
- If no firewood found thank the driver, provide EAB pamphlet, remove flagging, and direct back into traffic
- If illegal hardwood is present seize and safeguard, explain why firewood is illegal, complete survey record and a PPQ 523 EAN, take pictures of firewood with an identifier in the picture
- Inform the driver about penalty procedures

Other

Staff to relieve other positions (1-2)

Supervisor (minimum 1)

Total Staffing = 10 (numbers may be reduced if low volume roadway is selected)

#1 Priority – Safety

#1 Priority -- Courtesy

(Fill out for each traveler carrying regulated articles)

Inspector Name: _____ Agency: _____ Date: _____

Operation Stop Location: _____ Time: _____

<p>Identification and Origin of Individual LICENSE PLATE # IS MANDATORY</p>	<p>Name: _____ Address: _____ _____ City: _____ State: _____ Zip: _____ Phone: _____ License Plate (including state): _____</p>
<p>Individual/Regulated Item Destination (Campground, Business, etc.) COUNTY IS MANDATORY</p>	<p>Enter Destination Information here: COUNTY _____</p>
<p>Regulated Item Origin</p>	<p>Inside quarantine? _____ Yes _____ No Origin if different than traveler (county, address, etc.):</p>
<p>Traveler aware of EAB Quarantine/Regulations?</p>	<p>_____ No _____ Yes (if yes, how?) How? _____</p>
<p>Regulated Item Type: _____ Nursery stock _____ Logs/Lumber _____ Firewood _____ Other</p>	<p>If firewood mark blanks as appropriate: _____ Commercial Package _____ Non-commercial Ash present _____ Yes _____ Unknown</p>
<p>Amount of Regulated Item</p>	<p>_____ Piece Count _____ Estimated Weight</p>
<p>Regulatory Action</p>	<p>Seizure? _____ Yes _____ No If no why not? _____</p>
<p>Evidence of EAB –check those that apply→ EAB Inspection not conducted: _____</p>	<p>_____ D shaped exit holes _____ Bark splits _____ Serpentine galleries _____ EAB life stage</p>
<p>Comments:</p>	

Inspector Name: _____ Date: _____

Regulatory Contact

Concern or Business Name: _____

Contact Name: _____

Street Address: _____ City: _____

State: _____ Zip Code _____

County: _____

GPS Lat _____ Long _____

Phone: _____

Concern Type (circle up to three)

Nursery / Garden Center / Firewood Dealer / Public Campground / Private
Campground / Milling & Manufacturing / Landscaper / Research / Municipality /
Tree Removal Service / Landscaper / Logging Company / Misc. / Pallets / Pallet
Stock / Mulch

Aware of EAB Quarantine Y or N If yes,
how? _____

Interstate Movement Y or N

What/Commodity?

Frequency of shipments?

Quantity per shipment?

Final Destination of shipments?

From area near EAB positives?

Shipping Documents Available?

***Signature below confirms representative(s) of the Cooperative Emerald
Ash Borer Project contacted the signatory individual and gave them an
information packet about Emerald Ash Borer.***

Printed Name: _____

Signature: _____

Date: _____

Survey of Ash

Ash Product	Present? Y or N	Origin	Quantity	Inspected? Y or N	How long has the material been on site?
Firewood					
Nursery Stock					
Estab. Ash					
Ash Chips/Mulch					
Saw Logs					
Other					


Campground Survey

	Ash firewood Present? Y or N	% OH plates	% MI plates	% IN plates	Local Firewood Available	Permanent Campsites Present
Private Campground						
Public Campground						

EAB symptoms or specimens present Y or N,
 If yes record ash product and X all indicators that apply.
 Ash Product: _____

Crown Dieback	Life Stage	D-shaped holes	Bark splits	Epicormic shoots	Serpentine Gallery	Die-Back	Woodpecker Damage

Remarks:




Identification of Regulatory Concerns Useful Sources

Useful Sources to Identify Regulatory Concerns

A variety of methods are used by EAB regulatory staff to locate individuals/businesses that present a risk of spreading EAB. Below is a list of some of those that have been most effective along with some examples.

1. Physical Reconnaissance
2. Phone Directory Yellow Pages
 - a. Sources
 - Superpages.com
 - Switchboard.com
 - Yellowbook.com
 - Google.com
 - b. Keywords
 - Firewood
 - Lumber Wholesale
 - Sawmills & Planing mills
 - Campgrounds
 - Logging
3. Internet Auctions
 - a. Ebay
4. Trade magazines
 - a. Physical magazines
 - b. Internet newsletters
 - <http://www.timberlinemag.com/aboutTL.asp>
 - <http://www.timberpa.com/index2.htm>
 - <http://www.timberbuyer.net/forum.shtml>
 - <http://www.logsplitters.com/ArborAssociations.html>
5. Internet Chats/Blogs
 - a. Firewoodcenter.com
 - b. Firewood.com
6. Internet Search Engine
 - a. Google
 - b. Yahoo
 - c. MSN



United States Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine

Cooperative Emerald Ash Borer Project

Identification of Regulatory Concerns
Useful Sources

7. State records
 - a. LLC/Inc. registrations
8. County records
 - a. DBA (Doing Business As) Licenses
9. Forest Industry Directories
 - a. State Dept. Natural Resource
<http://www.michigandnr.com/wood/>
 - b. Forest Service
<http://www.srs.fs.usda.gov/econ/econhome.htm>
10. County Health Departments
Public Health - many states test private campground water sources
11. Personal Referrals
12. Targeted Operations

Appendix H

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information is 0579-0102. The time required to complete this information collection is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

FORM APPROVED - OMB NO. 0579-0102

U.S. DEPARTMENT OF AGRICULTURE ANIMAL AND PLANT HEALTH INSPECTION SERVICE PLANT PROTECTION AND QUARANTINE EMERGENCY ACTION NOTIFICATION	SERIAL NO. <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:50%;">1. PPQ LOCATION</td> <td style="width:50%;">2. DATE ISSUED</td> </tr> </table>	1. PPQ LOCATION	2. DATE ISSUED
1. PPQ LOCATION	2. DATE ISSUED		
3. NAME AND QUANTITY OF ARTICLE(S)	4. LOCATION OF ARTICLES		
6. SHIPPER	5. DESTINATION OF ARTICLES		
9. OWNER/CONSIGNEE OF ARTICLES	7. NAME OF CARRIER		
Name: _____ Address: _____ _____ _____ PHONE NO. _____ FAX NO. _____ SS NO. _____ TAX ID NO. _____	8. SHIPMENT ID NO.(S)		
	10. PORT OF LADING		
	11. DATE OF ARRIVAL		
	12. ID OF PEST(S), NOXIOUS WEEDS, OR ARTICLE(S)		
	12a. PEST ID NO.		
	12b. DATE INTERCEPTED		
	13. COUNTRY OF ORIGIN		
	14. GROWER NO.		
	15. FOREIGN CERTIFICATE NO.		
	15a. PLACE ISSUED		
	15b. DATE		

Under Sections 411, 412, and 414 of the Plant Protection Act (7 USC 7711, 7712, and 7714) and Sections 10404 through 10407 of the Animal Health Protection Act (7 USC 8303 through 8306), you are hereby notified, as owner or agent of the owner of said carrier, premises, and/or articles, to apply remedial measures for the pest(s), noxious weeds, and or article(s) specified in Item 12, in a manner satisfactory to and under the supervision of an Agriculture Officer. Remedial measures shall be in accordance with the action specified in Item 16 and shall be completed within the time specified in Item 17.

AFTER RECEIPT OF THIS NOTIFICATION, ARTICLES AND/OR CARRIERS HEREIN DESIGNATED MUST NOT BE MOVED EXCEPT AS DIRECTED BY AN AGRICULTURE OFFICER. THE LOCAL OFFICER MAY BE CONTACTED AT:

16. ACTION REQUIRED

- TREATMENT: _____
- RE-EXPORTATION: _____
- DESTRUCTION: _____
- OTHER: _____

Should the owner or owner's agent fail to comply with this order within the time specified below, USDA is authorized to recover from the owner or agent cost of any care, handling, application of remedial measures, disposal, or other action incurred in connection with the remedial action, destruction, or removal.

17. AFTER RECEIPT OF THIS NOTIFICATION COMPLETE SPECIFIED ACTION WITHIN (Specify No. Hours or No. Days):	18. SIGNATURE OF OFFICER:
--	---------------------------

ACKNOWLEDGMENT OF RECEIPT OF EMERGENCY ACTION NOTIFICATION

I hereby acknowledge receipt of the foregoing notification.

SIGNATURE AND TITLE:	DATE AND TIME:
----------------------	----------------

19. REVOCATION OF NOTIFICATION

ACTION TAKEN: _____

SIGNATURE OF OFFICER:	DATE:
-----------------------	-------

Appendix I

PPQ Form 530 – (Limited Permit)

No. C-325575

Information requested is needed to determine if a permit can be issued (7 CFR 301).
FORM APPROVED: OMB NOS. 0579-0008 & 0579-0123
See reverse side for additional information.

U.S. DEPARTMENT OF AGRICULTURE
ANIMAL AND PLANT HEALTH INSPECTION SERVICE
PLANT PROTECTION AND QUARANTINE

LIMITED PERMIT

This permit authorizes the movement of the NONCERTIFIED articles described below to a specified destination for limited handling, utilization, or processing, or for treatment. The movement of such articles is regulated by Federal or State cooperative domestic plant quarantines.

1. DATE ISSUED	2. VOID AFTER	
3. NAME OF CONSIGNOR		
4. SHIPPING POINT		
5. NAME AND ADDRESS OF CONSIGNEE		
6. VEHICLE LICENSE NO. & STATE		
7. R.R. CAR INITIALS		
II. DESCRIPTION		
A. Quantity	B. Article	C. Remarks
9. SIGNATURE OF ISSUING OFFICER		

ENDORSEMENT

The above described shipment was received by the designated consignee, and was handled in the manner approved under the provisions of all applicable Federal or State cooperative domestic plant quarantines.

10. DATE RECEIVED
11. SIGNATURE OF DESTINATION OFFICER

PENALTY FOR MISUSE OR ALTERATION (7 USC 163)

PPQ FORM 530 (FEB 2002)
Previous edition dated APR 58 may be used.

PART 1 - CONSIGNEE'S COPY

U.S. GOVERNMENT PRINTING OFFICE: 2002-722-410

Appendix J

PPQ Form 540 – (Certificate)

FORM APPROVED
OMB NO. 0579-0088

Information requested is needed to determine if a permit can be issued (7 CFR 301).
See reverse side for additional information.

089704

U.S. DEPARTMENT OF AGRICULTURE
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
CERTIFICATE
This certificate must be surrendered to the consignee at destination of shipment.

The articles described below are certified under all applicable Federal or State cooperative domestic plant quarantines.

1. DATE ISSUED _____ 2. VOID AFTER _____

3. NAME OF CONSIGNOR _____

4. SHIPPING POINT _____

5. NAME & ADDRESS OF CONSIGNEE _____

6. VEHICLE LICENSE NO. & STATE _____ 7. R.R. CAR INITIALS & NO. _____

8. DESCRIPTION

A. Quantity	B. Article	C. Remarks

9. SIGNATURE OF ISSUING INSPECTOR _____

PENALTY FOR MISUSE OR ALTERATION
(7 USC 163)

PPQ FORM 540 (APR 89) Previous edition obsolete. **PART 1—CONSIGNEE**

Passive dispersal pathways of emerald ash borer, *Agrilus planipennis* (Fairmaire), (Coleoptera: Buprestidae) identified by USDA APHIS PPQ EAB Program

Item: *Firewood*

Type: Packaged/bundled or bulk (split, unsplit, or slab)

Pathway: “Big box” store
 Firewood producer/distributor
 Firewood broker
 Sawmill slab wood
 Campground
 Online sale and auction
 Local auction
 Roadside sale
 Small business sale
 Homeowner collection

Method of transport: Long-distance hauling company
 Domestic parcel delivery service
 Private vehicle

Item: *Solid Wood Packing Material*

Type: Dunnage
 New and recycled pallet
 Pallet stock/cant
 Crating
 Case
 Skid
 Block

Pathway: Pallet manufacturer
 Sawmill
 Distribution center

Method of transport: Railway
 Freight
 Long-distance hauling company
 Domestic parcel delivery service
 Air cargo service

Item: *Lumber*

Type: Sawn timber
Green lumber
Air-dried rough lumber
Air-dried dimensioned lumber
Miscellaneous products
 Railroad ties
 Crane mats
 Stakes
 Trailer beds and sides

Pathway: Sawmill
Portable sawmill operator

Method of transport: Railway
Long-distance hauling company
Domestic parcel delivery service
Local delivery service

Item: *Logs*

Type: Saw log
Pulp log
Veneer log

Pathway: Logger
Log hauling company
Sawmill
Portable sawmill operator
Tree service company
Veneer mill
Paper mill
Pulp mill

Method of transport: Freightier/barge
Railway
Long-distance hauling company
Local hauling service

Item: *Chips*

Type: Biofuel
Animal bedding
Engineered wood panel material

Pathway: Wood-fired utility company
Panel manufacturer
Zoo
Tree service company
Sawmill

Method of transport: Long-distance hauling company
Railway
Local delivery service
Private vehicle

Item: *Mulch (Composted and uncomposted)*

Pathway: Landscape company

Method of transport: Long-distance hauling company
Local delivery service
Private vehicle

Item: *Nursery Stock (Fraxinus spp.)*

Pathway: Nursery
Online sales and auction

Method of transport: Long-distance hauling company
Local delivery service
Private vehicle